

**NMDA Receptor, NR2A Subunit N-terminus Antibody**  
Affinity purified rabbit polyclonal antibody  
Catalog # AN1150

### Specification

---

#### NMDA Receptor, NR2A Subunit N-terminus Antibody - Product Information

Application	WB, IHC
Primary Accession	<a href="#">Q00959</a>
Reactivity	Rat
Predicted	Bovine, Mouse
Host	Rabbit
Clonality	polyclonal
Calculated MW	180 KDa

#### NMDA Receptor, NR2A Subunit N-terminus Antibody - Additional Information

Gene ID	24409
Gene Name	GRIN2A

#### Other Names

Glutamate receptor ionotropic, NMDA 2A, GluN2A, Glutamate [NMDA] receptor subunit epsilon-1, N-methyl D-aspartate receptor subtype 2A, NMDAR2A, NR2A, Grin2a

#### Target/Specificity

Synthetic peptide corresponding to amino acid residues from the N-terminal region of the NR2A subunit conjugated to KLH.

#### Dilution

WB~~ 1:1000

IHC~~ 1:500

#### Format

Prepared from rabbit serum by affinity purification using a column to which the peptide immunogen was coupled.

#### Antibody Specificity

Specific for the ~180k NR2A subunit of the NMDA receptor.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

NMDA Receptor, NR2A Subunit N-terminus Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### Shipping


Blue Ice

## NMDA Receptor, NR2A Subunit N-terminus Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## NMDA Receptor, NR2A Subunit N-terminus Antibody - Images



Western blot of 10 ug of rat hippocampal lysate showing specific immunolabeling of the ~ 180k NR2A subunit.



Immunostaining of 21 DIV nucleofected mouse striatal neuron (green) co-cultured with cortical neurons showing nice punctate labeling of the N-terminal NR2A subunit (red) in both the medium spiny neurons and the large pyramidal cell in the upper right. Photo courtesy of Dr. A.J. Milnerwood, Dr. Lynn Raymond Lab, University of British Columbia.

## NMDA Receptor, NR2A Subunit N-terminus Antibody - Background

The ion channels activated by glutamate are typically divided into two classes. Glutamate receptors that are activated by kainate and  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxalone propionic acid (AMPA) are known as kainate/AMPA receptors (K/AMPA). Those that are sensitive to N-methyl-D-aspartate (NMDA) are designated NMDA receptors (NMDAR). The NMDAR plays an essential role in memory, neuronal development and it has also been implicated in several disorders of the central nervous system including Alzheimer's, epilepsy and ischemic neuronal cell

death (Grosshans et al., 2002; Wenthold et al., 2003; Carroll and Zukin, 2002). The NMDA receptor is also one of the principal molecular targets for alcohol in the CNS (Lovinger et al., 1989; Alvestad et al., 2003; Snell et al., 1996). The NMDAR is also potentiated by protein phosphorylation (Lu et al., 1999). The rat NMDAR1 (NR1) was the first subunit of the NMDAR to be cloned. The NR1 protein can form NMDA activated channels when expressed in *Xenopus* oocytes but the currents in such channels are much smaller than those seen in situ. Channels with more physiological characteristics are produced when the NR1 subunit is combined with one or more of the NMDAR2 (NR2 A-D) subunits.

### **NMDA Receptor, NR2A Subunit N-terminus Antibody - References**

- Alvestad RM, Grosshans DR, Coultrap SJ, Nakazawa T, Yamamoto T, Browning MD (2003) Tyrosine dephosphorylation and ethanol inhibition of N-methyl-D-aspartate receptor function. *J Biol Chem* 278:11020-11025.
- Carroll RC, Zukin RS (2002) NMDA-receptor trafficking and targeting: implications for synaptic transmission and plasticity. *Trends Neurosci* 25:571-577.
- Grosshans DR, Clayton DA, Coultrap SJ, Browning MD (2002) LTP leads to rapid surface expression of NMDA but not AMPA receptors in adult rat CA1. *Nat Neurosci* 5:27-33.
- Lovinger DM, White G, Weight FF (1989) Ethanol inhibits NMDA-activated ion current in hippocampal neurons. *Science* 243:1721-1724.
- Lu W-Y, Xiong Z-G, Lei S, Orser BA, Browning MD, MacDonald JF (1999) G-protein coupled receptors act via protein kinase C and Src to regulate NMDA receptors. *Nature Neurosci* 2:331-338.
- Snell LD, Nunley KR, Lickteig RL, Browning MD, Tabakoff B, Hoffman PL (1996) Regional and subunit specific changes in NMDA receptor mRNA and immunoreactivity in mouse brain following chronic ethanol ingestion. *Mol Brain Res* 40:71-78.
- Wenthold RJ, Prybylowski K, Standley S, Sans N, Petralia RS (2003) Trafficking of NMDA receptors. *Annu Rev Pharmacol Toxicol* 43:335-358.