

## Anti-COXIV (Ser58) Antibody

Our Anti-COXIV (Ser58) rabbit polyclonal phosphospecific primary antibody from PhosphoSolutions is p  
Catalog # AN1347

### Specification

---

#### Anti-COXIV (Ser58) Antibody - Product Information

Primary Accession	<a href="#">P19783</a>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>IgG</b>
Calculated MW	<b>19530</b>

#### Anti-COXIV (Ser58) Antibody - Additional Information

Gene ID **12857**

##### Other Names

AL024441 antibody, COX 4 antibody, COX IV 1 antibody, COX IV antibody, COX IV-1 antibody, Cox4 antibody, COX41\_HUMAN antibody, Cox4a antibody, COX4B antibody, COX4I1 antibody, COX4I2 antibody, COX4L2 antibody, Cytochrome c oxidase polypeptide IV antibody, Cytochrome c oxidase subunit 4 isoform 1 mitochondrial antibody, Cytochrome c oxidase subunit 4 isoform 1, mitochondrial antibody, Cytochrome C Oxidase subunit IV antibody, Cytochrome c oxidase subunit IV isoform 1 antibody, Cytochrome c oxidase subunit IV isoform 2 (lung) antibody, Cytochrome c oxydase subunit 4 antibody, MGC105470 antibody, MGC72016 antibody

##### Target/Specificity

COX, also known as cytochrome c oxidase, has 12 subunits that make up the transmembrane mitochondrial protein. Subunit IV has two isoforms; COXIV-1 and COXIV-2 (Huttermann et al., 2001). COXIV-1 is expressed ubiquitously while COXIV-2 is highly expressed in adult lung and low levels in brain and heart (Huttermann et al., 2001). Phosphorylation of amino acid residue Ser-58 of the COXIV-1 protein is a PKA-dependent regulation of COX and plays an important role in metabolism and CREB cycle activation (AcinPerez et al., 2011).

##### Format

Antigen Affinity Purified from Pooled Serum

##### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions

Anti-COXIV (Ser58) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

##### Shipping


Blue Ice

#### Anti-COXIV (Ser58) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Anti-COXIV (Ser58) Antibody - Images**



#### **Anti-COXIV (Ser58) Antibody - Background**

COX, also known as cytochrome c oxidase, has 12 subunits that make up the transmembrane mitochondrial protein. Subunit IV has two isoforms; COXIV-1 and COXIV-2 (Huttermann et al., 2001). COXIV-1 is expressed ubiquitously while COXIV-2 is highly expressed in adult lung and low levels in brain and heart (Huttermann et al., 2001). Phosphorylation of amino acid residue Ser-58 of the COXIV-1 protein is a PKA-dependent regulation of COX and plays an important role in metabolism and CREB cycle activation (AcinPerez et al., 2011).