

**Anti-CD46 (Extracellular region) Antibody**  
Catalog # AN1695**Specification****Anti-CD46 (Extracellular region) Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">P15529</a>
Host	Mouse
Clonality	Mouse Monoclonal
Isotype	IgG2b
Calculated MW	43747

**Anti-CD46 (Extracellular region) Antibody - Additional Information**

Gene ID 4179

**Other Names**

Membrane cofactor protein, TLX, Trophoblast leukocyte common antigen, CD46, MCP, MIC10

**Target/Specificity**

CD46 is a complement regulatory protein that is also called membrane cofactor protein. This protein is a type 1 membrane protein that plays an important inhibitory role in the complement system. CD46 exhibits a cofactor activity that promotes inactivation of C3b and C4b by serum factor 1, thereby protecting host cells from complement-dependent cytotoxicity. CD46 can also function as a receptor for selected bacteria and viruses, and is reportedly required for proper fusion of spermatozoa to the oocyte membrane during fertilization. CD46 is overexpressed in medulloblastoma tumors, and CD46 expression has been linked with poor prognosis in breast cancer. The upregulation of CD46 may protect cancer cells from complement-dependent cytotoxicity to facilitate cancer cell immune evasion.

**Format**

Protein G Purified

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Anti-CD46 (Extracellular region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Shipping**

Blue Ice


**Anti-CD46 (Extracellular region) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### Anti-CD46 (Extracellular region) Antibody - Images



Western blot of native lysates from NCI-H2052 mesothelial cells (lane 1), SK-MES-1 squamous carcinoma (lane 2), MDA-MB-231 breast carcinoma (lane 3), MeWo melanoma (lane 4), A431 skin adenocarcinoma (lane 5), LNCaP prostate cancer cells (lane 6), and MCF7 breast cancer cells (lane 7). The blot was probed with mouse monoclonal anti-CD46 (CM0371) at 1:1000.



Immunocytochemical labeling of CD46 in aldehyde fixed human NCIH1915 lung carcinoma cells. The cells were labeled with mouse monoclonal anti-CD46 (CM0371). The antibody was detected using goat anti-mouse Ig:DyLight® 594.



Representative Standard Curve using mouse monoclonal anti-CD46 (CM0371) for ELISA capture of human recombinant CD46 extracellular region with a His-tag. Captured protein was detected by suitable anti-His-tag antibody followed by appropriate secondary antibody HRP conjugate.

### **Anti-CD46 (Extracellular region) Antibody - Background**

CD46 is a complement regulatory protein that is also called membrane cofactor protein. This protein is a type 1 membrane protein that plays an important inhibitory role in the complement system. CD46 exhibits a cofactor activity that promotes inactivation of C3b and C4b by serum factor 1, thereby protecting host cells from complement-dependent cytotoxicity. CD46 can also function as a receptor for selected bacteria and viruses, and is reportedly required for proper fusion of spermatozoa to the oocyte membrane during fertilization. CD46 is overexpressed in medulloblastoma tumors, and CD46 expression has been linked with poor prognosis in breast cancer. The upregulation of CD46 may protect cancer cells from complement-dependent cytotoxicity to facilitate cancer cell immune evasion.