

**Anti-CD63 (Extracellular region) Antibody**  
Catalog # AN1701**Specification****Anti-CD63 (Extracellular region) Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P08962</a>
Host	Mouse
Clonality	Mouse Monoclonal
Isotype	IgG2a
Calculated MW	25637

**Anti-CD63 (Extracellular region) Antibody - Additional Information**

Gene ID 967

**Other Names**

CD63 antigen, Granulophysin, Lysosomal-associated membrane protein 3, LAMP-3, Melanoma-associated antigen ME491 OMA81H, Ocular melanoma-associated antigen, Tetraspanin-30, Tspan-30, MLA1, TSPAN30

**Target/Specificity**

Tetraspanins comprise a large superfamily of cell surface-associated membrane proteins with four transmembrane domains. On cell membranes, tetraspanins form networks of various proteins called tetraspanin-enriched microdomains (TEMs). CD63 was the first characterized tetraspanin and it is found in TEMs, as well as late endosomes and lysosomes. In late endosomes, CD63 is enriched on the intraluminal vesicles, and can be secreted as exosomes through fusion of endosomes with the plasma membrane. The complex localization pattern of CD63 suggests that its intracellular trafficking and distribution must be tightly regulated. CD63 contains N-link glycosylation sites that produce diverse CD63 molecules that range from 30 to 60 kDa depending on cell type. CD63 is an important exosomal marker in cancer cells, and may be involved in cancer progression and metastasis.

**Format**

Protein G Purified

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Anti-CD63 (Extracellular region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Shipping**


Blue Ice

**Anti-CD63 (Extracellular region) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Anti-CD63 (Extracellular region) Antibody - Images**



#### **Anti-CD63 (Extracellular region) Antibody - Background**

Tetraspanins comprise a large superfamily of cell surface-associated membrane proteins with four transmembrane domains. On cell membranes, tetraspanins form networks of various proteins called tetraspanin-enriched microdomains (TEMs). CD63 was the first characterized tetraspanin and it is found in TEMs, as well as late endosomes and lysosomes. In late endosomes, CD63 is enriched on the intraluminal vesicles, and can be secreted as exosomes through fusion of endosomes with the plasma membrane. The complex localization pattern of CD63 suggests that its intracellular trafficking and distribution must be tightly regulated. CD63 contains N-link glycosylation sites that produce diverse CD63 molecules that range from 30 to 60 kDa depending on cell type. CD63 is an important exosomal marker in cancer cells, and may be involved in cancer progression and metastasis.