

Anti-iNOS (Tyr-1055) [conserved site], Phosphospecific Antibody
Catalog # AN1869

Specification

Anti-iNOS (Tyr-1055) [conserved site], Phosphospecific Antibody - Product Information

Application	WB
Primary Accession	P35228
Reactivity	Bovine
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	131117

Anti-iNOS (Tyr-1055) [conserved site], Phosphospecific Antibody - Additional Information

Gene ID 4843

Other Names

Nos2 nitric oxide synthase 2, inducible, macrophage, NOS, type II, NOSII, Hepatocyte

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-iNOS (Tyr-1055) [conserved site], Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice


Anti-iNOS (Tyr-1055) [conserved site], Phosphospecific Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-iNOS (Tyr-1055) [conserved site], Phosphospecific Antibody - Images





Western blot analysis of mouse macrophages (J774A.1) treated with LPS (1 µg/ml) for 18 hrs followed by pervanadate (1 mM) for 30 min. (lanes 1, 3 & 5). The blots were then treated with alkaline phosphatase (lanes 2, 4 & 6). Blots were probed with rabbit polyclonal anti-inducible Nitric Oxide Synthase (iNOS) (lanes 1 & 2), anti-iNOS (Tyr-1055) (lanes 3 & 4), and mouse monoclonal anti-iNOS (lanes 5 & 6).



Immunocytochemical labeling of nNOS phosphorylation in rat PC12 cells differentiated with NGF. The cells were probed with mouse monoclonal (mAb) nNOS (NM4011), and rabbit polyclonal (pAb) nNOS (C-terminal region), nNOS (Tyr-895)/eNOS (Tyr-657), and nNOS (Tyr-1326)/iNOS (Tyr-1055). The antibodies were detected using appropriate secondary antibody conjugated to DyLight® 594.

Anti-iNOS (Tyr-1055) [conserved site], Phosphospecific Antibody - Background

Nitric oxide synthases (NOS), the enzymes responsible for synthesis of NO, are homodimers whose monomers are themselves two fused enzymes: a cytochrome reductase and a cytochrome that requires three cosubstrates (L-arginine, NADPH, and oxygen) and five cofactors or prosthetic groups (FAD, FMN, calmodulin, tetrahydrobiopterin, and heme). Several distinct NOS isoforms are produced from three distinct genes, inducible NOS (iNOS, NOS-II), neuronal NOS (nNOS, NOS-I), and endothelial NOS (eNOS, ecNOS, NOS-III). The inducible form of NOS, iNOS, is Ca²⁺ independent and is expressed in a broad range of cell types in response to stimulation with cytokines and exposure to microbial products. Phosphorylation of iNOS may regulate its activity and stability. Src kinase-induced phosphorylation of Tyr-1055 in iNOS reduces proteasomal degradation of iNOS, leading to increased NO production. Thus, phosphorylation may be an additional control for regulating iNOS activity in response to inflammatory conditions.