

## Anti-Osteopontin (N-terminal region) Antibody Catalog # AN1873

### Specification

---

#### Anti-Osteopontin (N-terminal region) Antibody - Product Information

Application	WB
Primary Accession	<a href="#">P10451</a>
Reactivity	Bovine
Host	Mouse
Clonality	Mouse Monoclonal
Isotype	IgG1
Calculated MW	35423

#### Anti-Osteopontin (N-terminal region) Antibody - Additional Information

Gene ID **6696**

##### Other Names

Bone sialoprotein 1, Nephropontin, Secreted phosphoprotein 1, SSP-1, Urinary stone protein, Uropontin, BNSP, OPN, SSP1, osteopontin

##### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions

Anti-Osteopontin (N-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

##### Shipping


Blue Ice


#### Anti-Osteopontin (N-terminal region) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-Osteopontin (N-terminal region) Antibody - Images





Western blot analysis of human recombinant osteopontin protein (lane 1 & 2) or human MDA-MB-231 cells (lane 3 & 4). The blots were probed with mouse monoclonal anti-Osteopontin (OM5741) at a dilution of 1:1000 (lane 1), 1:4000 (lane 2), 1:250 (lane 3), and 1:1000 (lane 4).



Immunocytochemical labeling of Osteopontin in paraformaldehyde fixed and NP-40 permeabilized MCF-7 cells. The cells were labeled with mouse monoclonal anti-Osteopontin (M574). The antibody was detected using goat anti-mouse DyLight® 594.

### Anti-Osteopontin (N-terminal region) Antibody - Background

Osteopontin (OPN) is a 34 kDa sialic acid rich member of small integrin-binding ligand N-linked glycoproteins. It is expressed in many different tissues and is post-translationally modified at multiple sites with both glycosylation and phosphorylation. The mature post-translationally modified protein is 60 kDa. OPN is involved with cell survival, proliferation, invasion, and stem like behavior. OPN can interact with CD44, bind hydroxyapatite, and activates many integrins. These interactions are important for OPN function in cell matrix formation. A higher presence of OPN has been found in a variety of cancers, leading to increased tumor growth and metastasis. In addition, OPN is involved in type I immunity through its function as a cytokine where it can enhance production of interferon-gamma and interleukin-12 and reduce production of interleukin-10.