

## Anti-RCAN1/Dscr1 (central region) Antibody Catalog # AN1936

### Specification

---

#### Anti-RCAN1/Dscr1 (central region) Antibody - Product Information

Application	WB
Primary Accession	<a href="#">P53805</a>
Reactivity	Bovine
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	28079

#### Anti-RCAN1/Dscr1 (central region) Antibody - Additional Information

Gene ID 1827

##### Other Names

Dscr1, MCIP, RCAN1, calcipressin, Adapt78

##### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions

Anti-RCAN1/Dscr1 (central region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

##### Shipping


Blue Ice


#### Anti-RCAN1/Dscr1 (central region) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-RCAN1/Dscr1 (central region) Antibody - Images





Western blot analysis of RCAN1 expression in human Jurkat (lanes 1 & 4), mouse C2C12 (lanes 2 & 5), and rat PC12 (lanes 3 & 6). The blot was probed with rabbit polyclonal anti-RCAN1 (C-terminus) at 1:1000 and anti-RCAN1 (a.a. 132-140) at 1:500.



### Anti-RCAN1/Dscr1 (central region) Antibody - Background

An important element of calcium signaling pathways involves calmodulin activation of calcineurin (phosphatase PP2B), leading to dephosphorylation of transcription factors such as NFAT and MEF2. A wide variety of proteins other than calmodulin have also been implicated in regulating calcineurin activity. Regulators of Calcineurin (RCANs) include RCAN1, RCAN2, and RCAN3. RCAN1 has previously been referred to as Down's syndrome candidate region-1 (Dscr1), MCIP, calcipressin, and Adapt78. This RCAN is expressed as several different variants with RCAN1L (38 kDa) and RCAN1S (31 kDa) being most prevalent. RCAN1 is increased in Down's syndrome tissues and in a mouse model of Down's syndrome. Increased expression of RCAN1 leads to significant suppression of tumor growth in mice as result of deficits in calcineurin-induced tumor angiogenesis. RCAN1 can recruit TAB1, TAK1, and calcineurin into a macromolecular signaling complex, and TAK1 can phosphorylate Ser-94 and Ser-136 in RCAN1S. This phosphorylation converts RCAN1 from an inhibitor to a facilitator of calcineurin-NFAT signaling.