

Anti-Stat5 (Tyr-694), Phosphospecific Antibody Catalog # AN1981

Specification

Anti-Stat5 (Tyr-694), Phosphospecific Antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | P42229 |
| Reactivity | Bovine |
| Host | Mouse |
| Clonality | Mouse Monoclonal |
| Isotype | IgG1 |
| Calculated MW | 90647 |

Anti-Stat5 (Tyr-694), Phosphospecific Antibody - Additional Information

| | |
|---|------|
| Gene ID | 6776 |
| Other Names | |
| Signal transducer and activator of transcription 5A, Stat5A | |

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Stat5 (Tyr-694), Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice


Anti-Stat5 (Tyr-694), Phosphospecific Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Stat5 (Tyr-694), Phosphospecific Antibody - Images





Western blot analysis of human A431 cells untreated (lanes 1, 3, 5, 7, & 9) or treated with EGF (100 nM) for 60 min (lanes 2, 4, 6, 8, & 10). The blots were probed with anti-Stat1 (lanes 1 & 2), anti-Stat1 (Tyr-701) (lanes 3 & 4), anti-Stat3 (lanes 5 & 6), anti-Stat5 (lanes 7 & 8), and anti-Stat5 (Tyr-694) (lanes 9 & 10).



Immunocytochemical labeling of Stat5 in control and pervanadate-treated A431 cells. The cells were labeled with mouse monoclonal Stat5 (SM2511) or Stat5 (Tyr-694) (SM1481) antibodies, then the antibodies were detected using appropriate secondary antibody conjugated to DyLight® 594.

Anti-Stat5 (Tyr-694), Phosphospecific Antibody - Background

The stat proteins function both as cytoplasmic signal transducers and as activators of transcription. Stat5 is activated in response to a wide variety of ligands including IL-2, GM-CSF, growth hormone, and prolactin. Phosphorylation at Tyr-694 is required for Stat5A activation. Stat5 has been found to be constitutively active in some leukemic cell types. Phosphorylated Stat5 is found in some endothelial cells treated with IL-3, which suggests its involvement in angiogenesis and cell motility. Both Stat5A (Tyr-694) and Stat5B (Tyr-699) are independently regulated and activated in various cell types. For instance, both isoforms are activated in response to IFN α in B cells, but only Stat5A is phosphorylated in response to IFN α in HeLa cells.