

Anti-Vav2 Antibody Catalog # AN2011

Specification

Anti-Vav2 Antibody - Product Information

Primary Accession	P52735
Reactivity	Bovine, Chicken
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	101289

Anti-Vav2 Antibody - Additional Information

Gene ID **7410**

Other Names

VAV2, Guanine nucleotide exchange factor VAV2

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Vav2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping

Blue Ice


Anti-Vav2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Vav2 Antibody - Images





Western blot of human Jurkat (lanes 1 & 4), HUVEC (lanes 2 & 5), and A431 (lanes 3 & 6) cells. The blots were probed with anti-Vav2 (a.a. 309-322) at a dilution of 1:500 (lanes 1-3) and anti-Vav3 (a.a. 293-305) at 1:500 (lanes 4-6).

Anti-Vav2 Antibody - Background

The Vav family of Rho-guanine nucleotide exchange factors, Vav1, Vav2, and Vav3, have central roles in transducing signals from cell surface receptors, such as integrin, growth factor and immune cell receptors to the cytoskeleton. This role includes receptor-mediated changes in the actin cytoskeleton and cell motility. Vav1 expression is normally restricted to hematopoietic cells, while Vav2 and Vav3 are more widely expressed. All three Vav isoforms have been shown to be abnormally expressed in several types of cancer. Vavs are composed of multiple domains, including a Dbl homology domain, a calponin homology domain, an acidic amino acid region, a pleckstrin homology domain, a cysteine-rich domain, and SH3 and SH2 domains. Vav activity is regulated by the phosphorylation status of several conserved tyrosine residues in the acidic region (In Vav2: Tyr-142, Tyr-159, and Tyr-172). These tyrosine residues are able to participate in autoinhibitory interactions with the Dbl homology domain and this interaction is prevented after phosphorylation of these sites leading to activation of Vav GEF activity.