

Anti-Ku80 Antibody
Catalog # AN2146**Specification**

Anti-Ku80 Antibody - Product Information

Primary Accession	P13010
Reactivity	Bovine, Chicken
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	82705

Anti-Ku80 Antibody - Additional InformationGene ID **7520****Other Names**

XRCC5, X-ray repair cross-complementing protein 5, ATP-dependent DNA helicase 2 subunit 2, Ku86, CTC box-binding factor 85 kDa subunit, Lupus Ku autoantigen protein p86, Thyroid-lupus autoantigen, TLAA, G22P2

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Ku80 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping

Blue Ice

Anti-Ku80 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Ku80 Antibody - Images**Anti-Ku80 Antibody - Background**

Telomere length maintenance, an activity essential for chromosome stability and genome integrity,

is regulated by telomerase- and telomere-associated factors. The DNA repair protein Ku (a heterodimer of Ku70 and Ku80 subunits) associates with mammalian telomeres and contributes to telomere maintenance. The Ku heterodimer functions at two kinds of DNA ends: telomeres and double-strand breaks. The role that Ku plays at these two classes of termini must be distinct, because Ku is required for accurate and efficient joining of double-strand breaks while similar DNA repair events are normally prohibited at chromosome ends