

Tyk2 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1072a**Specification**

Tyk2 Antibody - Product Information

Application	WB, IHC
Primary Accession	P29597
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1

Description

Tyk2 (tyrosine kinase 2), with 1187-amino acid protein (about 131kDa), belongs to the family of non-receptor janus tyrosine kinases, which also includes Jak1, Jak2, and Jak3. Kinases of the Jak family regulate a spectrum of cellular functions downstream of activated cytokine receptors in the lympho-hematopoietic system. Tyk2 is activated by a variety of cytokines: IFN-alpha, IFN-beta, IL-6, IL-10, IL-12, and IL-13 and promotes IFN-gamma production by Th1-type CD4 cells. Tyk2 can be viewed as a dual-function Jak, mediating both pro-inflammatory and anti-inflammatory cytokine responses. Tyk2 is also an important regulator of lymphoid tumor surveillance.

Immunogen

Purified recombinant fragment of Tyk2 expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

Tyk2 Antibody - Additional Information

Gene ID 7297

Other Names

Non-receptor tyrosine-protein kinase TYK2, 2.7.10.2, TYK2

Dilution

WB~~1/500 - 1/2000

IHC~~1:200~~1000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Tyk2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Tyk2 Antibody - Protein Information

Name TYK2

Function

Tyrosine kinase of the non-receptor type involved in numerous cytokines and interferons signaling, which regulates cell growth, development, cell migration, innate and adaptive immunity (PubMed:10542297, PubMed:10995743, PubMed:7657660, PubMed:7813427, PubMed:8232552). Plays both structural and catalytic roles in numerous interleukins and interferons (IFN-alpha/beta) signaling (PubMed:10542297). Associates with heterodimeric cytokine receptor complexes and activates STAT family members including STAT1, STAT3, STAT4 or STAT6 (PubMed:10542297, PubMed:7638186). The heterodimeric cytokine receptor complexes are composed of (1) a TYK2-associated receptor chain (IFNAR1, IL12RB1, IL10RB or IL13RA1), and (2) a second receptor chain associated either with JAK1 or JAK2 (PubMed:10542297, PubMed:25762719, PubMed:7526154, PubMed:7813427). In response to cytokine-binding to receptors, phosphorylates and activates receptors (IFNAR1, IL12RB1, IL10RB or IL13RA1), creating docking sites for STAT members (PubMed:7526154, PubMed:7657660). In turn, recruited STATs are phosphorylated by TYK2 (or JAK1/JAK2 on the second receptor chain), form homo- and heterodimers, translocate to the nucleus, and regulate cytokine/growth factor responsive genes (PubMed:10542297, PubMed:25762719, PubMed:7657660). Negatively regulates STAT3 activity by promoting phosphorylation at a specific tyrosine that differs from the site used for signaling (PubMed:29162862).

Tissue Location

Observed in all cell lines analyzed. Expressed in a variety of lymphoid and non-lymphoid cell lines

Tyk2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Tyk2 Antibody - Images






Figure 1: Western blot analysis using TYK2 mouse mAb against truncated TYK2 recombinant protein (1) and Jurkat cell lysate(2).




Figure 1: Immunohistochemical analysis of paraffin-embedded human breast tissue (A), lymph tissue (B) and skin carcinoma (C), showing membrane localization using BLK mouse mAb with DAB staining.

Tyk2 Antibody - References

1. Michael H. Shaw, Gordon J. Freeman, Mark F. Scott. *J. Immunol.*, Jun 2006; 176: 7263-7271.
2. Yohei Seto, Hiroshi Nakajima, Akira Suto. *J. Immunol.*, Jan 2003; 170: 1077.