

**PTK6 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1096a****Specification**

---

**PTK6 Antibody - Product Information**

Application	<b>WB, IHC</b>
Primary Accession	<a href="#">O13882</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG2b</b>
Calculated MW	<b>52kDa KDa</b>

**Description**

PTK6 (protein tyrosine kinase 6, BRK or FLJ42088), with 451-amino acid protein (about 52kDa), encodes a cytoplasmic nonreceptor protein kinase which may function as an intracellular signal transducer in epithelial tissues. Its presence in the nucleus appears to be linked to suppression of tumor progression. The encoded protein has been shown to undergo autophosphorylation. Very high level in colon and high levels in small intestine and prostate, and low levels in some fetal tissues. And Expressed at low level in some breast tumors, but not in normal breast. Also found in melanocytes, but not expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Overexpression of this gene in mammary epithelial cells leads to sensitization of the cells to epidermal growth factor and results in a partially transformed phenotype.

**Immunogen**

Purified recombinant fragment of human PTK6 expressed in E. Coli. <br /> <br />

**Formulation**

Ascitic fluid containing 0.03% sodium azide.

**PTK6 Antibody - Additional Information**

**Gene ID** 5753

**Other Names**

Protein-tyrosine kinase 6, 2.7.10.2, Breast tumor kinase, Tyrosine-protein kinase BRK, PTK6, BRK

**Dilution**

WB~~1/500 - 1/2000

IHC~~1:200~~1000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PTK6 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## PTK6 Antibody - Protein Information

**Name** PTK6

**Synonyms** BRK

### Function

Non-receptor tyrosine-protein kinase implicated in the regulation of a variety of signaling pathways that control the differentiation and maintenance of normal epithelia, as well as tumor growth. Function seems to be context dependent and differ depending on cell type, as well as its intracellular localization. A number of potential nuclear and cytoplasmic substrates have been identified. These include the RNA-binding proteins: KHDRBS1/SAM68, KHDRBS2/SLM1, KHDRBS3/SLM2 and SFPQ/PSF; transcription factors: STAT3 and STAT5A/B and a variety of signaling molecules: ARHGAP35/p190RhoGAP, PXN/paxillin, BTK/ATK, STAP2/BKS. Associates also with a variety of proteins that are likely upstream of PTK6 in various signaling pathways, or for which PTK6 may play an adapter-like role. These proteins include ADAM15, EGFR, ERBB2, ERBB3 and IRS4. In normal or non- tumorigenic tissues, PTK6 promotes cellular differentiation and apoptosis. In tumors PTK6 contributes to cancer progression by sensitizing cells to mitogenic signals and enhancing proliferation, anchorage-independent survival and migration/invasion. Association with EGFR, ERBB2, ERBB3 may contribute to mammary tumor development and growth through enhancement of EGF-induced signaling via BTK/AKT and PI3 kinase. Contributes to migration and proliferation by contributing to EGF-mediated phosphorylation of ARHGAP35/p190RhoGAP, which promotes association with RASA1/p120RasGAP, inactivating RhoA while activating RAS. EGF stimulation resulted in phosphorylation of PNXPaxillin by PTK6 and activation of RAC1 via CRK/CrKII, thereby promoting migration and invasion. PTK6 activates STAT3 and STAT5B to promote proliferation. Nuclear PTK6 may be important for regulating growth in normal epithelia, while cytoplasmic PTK6 might activate oncogenic signaling pathways.

### Cellular Location

Cytoplasm. Nucleus. Cell projection, ruffle. Membrane. Note=Colocalizes with KHDRBS1, KHDRBS2 or KHDRBS3, within the nucleus. Nuclear localization in epithelial cells of normal prostate but cytoplasmic localization in cancer prostate

### Tissue Location

Epithelia-specific. Very high level in colon and high levels in small intestine and prostate, and low levels in some fetal tissues. Not expressed in breast or ovarian tissue but expressed in high percentage of breast and ovarian cancers. Also overexpressed in some metastatic melanomas, lymphomas, colon cancers, squamous cell carcinomas and prostate cancers. Also found in melanocytes. Not expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Isoform 2 is present in prostate epithelial cell lines derived from normal prostate and prostate adenocarcinomas, as well as in a variety of cell lines.

## PTK6 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## PTK6 Antibody - Images




Figure 1: Western blot analysis using PTK6 mouse mAb against HeLa (1), A549 (2) and MCF-7 (3) cell lysate.




Figure 2: Immunohistochemical analysis of paraffin-embedded malignant mesenchymoma tissues, showing cytoplasmic localization using C-kit mouse mAb with DAB staining.

#### PTK6 Antibody - References

1. Wilks, A.F. Proc. Natl. Acad. Sci. USA 86: 1603-1607.
2. Otilie, S., et al. Oncogene 7: 1625-1630.
3. Lee, S.T., et al. Oncogene 8: 3403-3410.