

**NCOA3 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1225a****Specification**

---

**NCOA3 Antibody - Product Information**

Application	<b>WB, FC, IHC</b>
Primary Accession	<a href="#">O9Y6Q9</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG1</b>

**Description**

NCOA3: nuclear receptor coactivator 3. The protein encoded by this gene is a nuclear receptor coactivator that interacts with nuclear hormone receptors to enhance their transcriptional activator functions. The encoded protein has histone acetyltransferase activity and recruits p300/CBP-associated factor and CREB binding protein as part of a multisubunit coactivation complex. This protein is initially found in the cytoplasm but is translocated into the nucleus upon phosphorylation. Two transcript variants encoding different isoforms have been found for this gene. In addition, a polymorphic repeat region is found in the C-terminus of the encoded protein.

**Immunogen**

Purified recombinant fragment of NCOA3 (aa1-200) expressed in E. Coli. <br />

**Formulation**

Ascitic fluid containing 0.03% sodium azide. <br />

**NCOA3 Antibody - Additional Information**

**Gene ID** 8202

**Other Names**

Nuclear receptor coactivator 3, NCoA-3, 2.3.1.48, ACTR, Amplified in breast cancer 1 protein, AIB-1, CBP-interacting protein, pCIP, Class E basic helix-loop-helix protein 42, bHLHe42, Receptor-associated coactivator 3, RAC-3, Steroid receptor coactivator protein 3, SRC-3, Thyroid hormone receptor activator molecule 1, TRAM-1, NCOA3, AIB1, BHLHE42, RAC3, TRAM1

**Dilution**

WB~~1/500 - 1/2000

FC~~1:200~~400

IHC~~1:200~~1000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

NCOA3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## NCOA3 Antibody - Protein Information

**Name** NCOA3

**Synonyms** AIB1, BHLHE42, RAC3, TRAM1

### Function

Nuclear receptor coactivator that directly binds nuclear receptors and stimulates the transcriptional activities in a hormone- dependent fashion. Plays a central role in creating a multisubunit coactivator complex, which probably acts via remodeling of chromatin. Involved in the coactivation of different nuclear receptors, such as for steroids (GR and ER), retinoids (RARs and RXRs), thyroid hormone (TRs), vitamin D3 (VDR) and prostanoids (PPARs). Displays histone acetyltransferase activity. Also involved in the coactivation of the NF-kappa-B pathway via its interaction with the NFKB1 subunit.

### Cellular Location

Cytoplasm. Nucleus. Note=Mainly cytoplasmic and weakly nuclear. Upon TNF activation and subsequent phosphorylation, it translocates from the cytoplasm to the nucleus

### Tissue Location


Widely expressed. High expression in heart, skeletal muscle, pancreas and placenta. Low expression in brain, and very low in lung, liver and kidney

## NCOA3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## NCOA3 Antibody - Images






Figure 1: Western blot analysis using NCOA3 mouse mAb against truncated Trx-NCOA3 recombinant protein (1) and truncated NCOA3 (aa1-200)-hlgGfC transfected CHOK1 cell lysate (2).




Figure 7: Flow cytometric analysis of HeLa cells using CD44 mouse mAb (right) and negative control (left).




Figure 3: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissues, showing membrane localization with DAB staining using CD44 mouse mAb.

**NCOA3 Antibody - References**

1. Mol Cell Biol. 2005 Sep;25(18):8273-84.
2. J Clin Oncol. 2006 Oct 1;24(28):4565-9.