

AGT Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1300a**Specification**

AGT Antibody - Product Information

Application	WB, FC, ICC, IF
Primary Accession	P01019
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Calculated MW	53kDa KDa

Description

AGT, pre-angiotensinogen or angiotensinogen precursor, is expressed in the liver and is cleaved by the enzyme renin in response to lowered blood pressure. The resulting product, angiotensin I, is then cleaved by angiotensin converting enzyme (ACE) to generate the physiologically active enzyme angiotensin II. The protein is involved in maintaining blood pressure and in the pathogenesis of essential hypertension and preeclampsia. Mutations in this gene are associated with susceptibility to essential hypertension, and can cause renal tubular dysgenesis, a severe disorder of renal tubular development. Defects in this gene have also been associated with non-familial structural atrial fibrillation, and inflammatory bowel disease.

Immunogen

Purified recombinant fragment of human AGT expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

AGT Antibody - Additional Information

Gene ID 183

Other Names

Angiotensinogen, Serpin A8, Angiotensin-1, Angiotensin 1-10, Angiotensin I, Ang I, Angiotensin-2, Angiotensin 1-8, Angiotensin II, Ang II, Angiotensin-3, Angiotensin 2-8, Angiotensin III, Ang III, Des-Asp[1]-angiotensin II, Angiotensin-4, Angiotensin 3-8, Angiotensin IV, Ang IV, Angiotensin 1-9, Angiotensin 1-7, Angiotensin 1-5, Angiotensin 1-4, AGT, SERPINA8

Dilution

WB~~1/500 - 1/2000
FC~~1:200~~400
ICC~~1:200~~1000
IF~~1:200~1000.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

AGT Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

AGT Antibody - Protein Information

Name AGT ([HGNC:333](#))

Synonyms SERPINA8

Function

Essential component of the renin-angiotensin system (RAS), a potent regulator of blood pressure, body fluid and electrolyte homeostasis.

Cellular Location

Secreted

Tissue Location

Expressed by the liver and secreted in plasma.

AGT Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

AGT Antibody - Images




Figure 1: Western blot analysis using AGT mouse mAb against human plasma (1).




Figure 3: Flow cytometric analysis of A549 cells using TUBB3 mouse mAb (green) and negative control (purple).




Figure 2: Immunofluorescence analysis of PANC-1 cells using TUBB3 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 2: Immunofluorescence analysis of PANC-1 cells using TUBB3 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.

AGT Antibody - References

1. Am J Physiol Heart Circ Physiol. 2007 Sep;293(3):H1900-7.
2. Regul Pept. 2006 Jan 15;133(1-3):155-9.