

BMP4 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1340a**Specification**

BMP4 Antibody - Product Information

Application	WB, ICC, IF
Primary Accession	P12644
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	64kDa KDa

Description

The protein encoded by this gene is a member of the bone morphogenetic protein family which is part of the transforming growth factor-beta superfamily. The superfamily includes large families of growth and differentiation factors. BMPs (bone morphogenetic proteins) belong to the TGF beta superfamily of structurally related signaling proteins. Members of this superfamily are widely represented throughout the animal kingdom and have been implicated in a variety of developmental processes. Proteins of the TGF beta superfamily are disulfide-linked dimers composed of two 12-15 kDa polypeptide chains. As implied by their name, BMPs initiate, promote and regulate bone development, growth, remodeling and repair. Smad1 translocation to the nucleus is observed after the addition of BMP4 (also designated BMP2B), suggesting that BMP4 may play a role in activation of the Smad pathway. BMP is secreted into the extracellular matrix.

Immunogen

Purified recombinant fragment of human BMP4 expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

BMP4 Antibody - Additional Information

Gene ID 652

Other Names

Bone morphogenetic protein 4, BMP-4, Bone morphogenetic protein 2B, BMP-2B, BMP4, BMP2B, DVR4

Dilution

WB~~1/500 - 1/2000
ICC~~1:200~~1000
IF~~1:200~1000.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BMP4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

BMP4 Antibody - Protein Information

Name BMP4 ([HGNC:1071](#))

Function

Growth factor of the TGF-beta superfamily that plays essential roles in many developmental processes, including neurogenesis, vascular development, angiogenesis and osteogenesis (PubMed: [31363885](http://www.uniprot.org/citations/31363885)). Acts in concert with PTHLH/PTHRP to stimulate ductal outgrowth during embryonic mammary development and to inhibit hair follicle induction (By similarity). Initiates the canonical BMP signaling cascade by associating with type I receptor BMPRI1A and type II receptor BMPRI2 (PubMed: [25868050](http://www.uniprot.org/citations/25868050), PubMed: [8006002](http://www.uniprot.org/citations/8006002)). Once all three components are bound together in a complex at the cell surface, BMPRI2 phosphorylates and activates BMPRI1A. In turn, BMPRI1A propagates signal by phosphorylating SMAD1/5/8 that travel to the nucleus and act as activators and repressors of transcription of target genes (PubMed: [25868050](http://www.uniprot.org/citations/25868050), PubMed: [29212066](http://www.uniprot.org/citations/29212066)). Positively regulates the expression of odontogenic development regulator MSX1 via inducing the IPO7- mediated import of SMAD1 to the nucleus (By similarity). Required for MSX1-mediated mesenchymal molar tooth bud development beyond the bud stage, via promoting Wnt signaling (By similarity). Acts as a positive regulator of odontoblast differentiation during mesenchymal tooth germ formation, expression is repressed during the bell stage by MSX1- mediated inhibition of CTNNA1 signaling (By similarity). Able to induce its own expression in dental mesenchymal cells and also in the neighboring dental epithelial cells via an MSX1-mediated pathway (By similarity). Can also signal through non-canonical BMP pathways such as ERK/MAP kinase, PI3K/Akt, or SRC cascades (PubMed: [31363885](http://www.uniprot.org/citations/31363885)). For example, induces SRC phosphorylation which, in turn, activates VEGFR2, leading to an angiogenic response (PubMed: [31363885](http://www.uniprot.org/citations/31363885)).

Cellular Location

Secreted, extracellular space, extracellular matrix

Tissue Location

Expressed in the lung and lower levels seen in the kidney. Present also in normal and neoplastic prostate tissues, and prostate cancer cell lines

BMP4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BMP4 Antibody - Images




Figure 1: Western blot analysis using BMP4 mouse mAb against BMP4-hlgGfC transfected HEK293 cell lysate.




Figure 2: Immunofluorescence analysis of PC-3 (left) and SK-BR-3 (right) cells using anti-GOT2 mAb (green). Red: Actin filaments have been labeled with DY-554 phalloidin. Blue: DRAQ5 fluorescent DNA dye.




Figure 2: Immunofluorescence analysis of PC-3 (left) and SK-BR-3 (right) cells using anti-GOT2 mAb (green). Red: Actin filaments have been labeled with DY-554 phalloidin. Blue: DRAQ5 fluorescent DNA dye.

BMP4 Antibody - References

1. Genomics. 1995 Jun 10;27(3):559-60.
2. DNA Seq. 1995;5(5):273-5.
3. J Bone Miner Res. 2009 Dec;24(12):2039-49.
4. Stem Cells Dev. 2009 Nov;18(9):1283-92.