

CHUK Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1458a

Specification

CHUK Antibody - Product Information

Application	WB, IF, FC
Primary Accession	O15111
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	85kDa KDa

Description

This gene encodes a member of the serine/threonine protein kinase family. The encoded protein, a component of a cytokine-activated protein complex that is an inhibitor of the essential transcription factor NF-kappa-B complex, phosphorylates sites that trigger the degradation of the inhibitor via the ubiquitination pathway, thereby activating the transcription factor.

Immunogen

Purified recombinant fragment of human CHUK expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

CHUK Antibody - Additional Information

Gene ID 1147

Other Names

Inhibitor of nuclear factor kappa-B kinase subunit alpha, I-kappa-B kinase alpha, IKK-A, IKK-alpha, Ikbka, IkappaB kinase, 2.7.11.10, Conserved helix-loop-helix ubiquitous kinase, I-kappa-B kinase 1, IKK1, Nuclear factor NF-kappa-B inhibitor kinase alpha, NFKBIKA, Transcription factor 16, TCF-16, CHUK, IKKA, TCF16

Dilution

WB~~1/500 - 1/2000

IF~~1/200 - 1/1000

FC~~1/200 - 1/400

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CHUK Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CHUK Antibody - Protein Information

Name CHUK

Synonyms IKKA, TCF16

Function

Serine kinase that plays an essential role in the NF-kappa-B signaling pathway which is activated by multiple stimuli such as inflammatory cytokines, bacterial or viral products, DNA damages or other cellular stresses (PubMed: [18626576](http://www.uniprot.org/citations/18626576)), PubMed: [9244310](http://www.uniprot.org/citations/9244310)), PubMed: [9252186](http://www.uniprot.org/citations/9252186)), PubMed: [9346484](http://www.uniprot.org/citations/9346484)). Acts as a part of the canonical IKK complex in the conventional pathway of NF-kappa-B activation and phosphorylates inhibitors of NF-kappa-B on serine residues (PubMed: [18626576](http://www.uniprot.org/citations/18626576)), PubMed: [35952808](http://www.uniprot.org/citations/35952808)), PubMed: [9244310](http://www.uniprot.org/citations/9244310)), PubMed: [9252186](http://www.uniprot.org/citations/9252186)), PubMed: [9346484](http://www.uniprot.org/citations/9346484)). These modifications allow polyubiquitination of the inhibitors and subsequent degradation by the proteasome (PubMed: [18626576](http://www.uniprot.org/citations/18626576)), PubMed: [9244310](http://www.uniprot.org/citations/9244310)), PubMed: [9252186](http://www.uniprot.org/citations/9252186)), PubMed: [9346484](http://www.uniprot.org/citations/9346484)). In turn, free NF-kappa-B is translocated into the nucleus and activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis (PubMed: [18626576](http://www.uniprot.org/citations/18626576)), PubMed: [9244310](http://www.uniprot.org/citations/9244310)), PubMed: [9252186](http://www.uniprot.org/citations/9252186)), PubMed: [9346484](http://www.uniprot.org/citations/9346484)). Negatively regulates the pathway by phosphorylating the scaffold protein TAXBP1 and thus promoting the assembly of the A20/TNFAIP3 ubiquitin-editing complex (composed of A20/TNFAIP3, TAX1BP1, and the E3 ligases ITCH and RNF11) (PubMed: [21765415](http://www.uniprot.org/citations/21765415)). Therefore, CHUK plays a key role in the negative feedback of NF-kappa-B canonical signaling to limit inflammatory gene activation. As part of the non-canonical pathway of NF-kappa-B activation, the MAP3K14-activated CHUK/IKKA homodimer phosphorylates NFKB2/p100 associated with RelB, inducing its proteolytic processing to NFKB2/p52 and the formation of NF-kappa-B RelB-p52 complexes (PubMed: [20501937](http://www.uniprot.org/citations/20501937)). In turn, these complexes regulate genes encoding molecules involved in B-cell survival and lymphoid organogenesis. Participates also in the negative feedback of the non-canonical NF- kappa-B signaling pathway by phosphorylating and destabilizing MAP3K14/NIK. Within the nucleus, phosphorylates CREBBP and consequently increases both its transcriptional and histone acetyltransferase activities (PubMed: [17434128](http://www.uniprot.org/citations/17434128)). Modulates chromatin accessibility at NF- kappa-B-responsive promoters by phosphorylating histones H3 at 'Ser-10' that are subsequently acetylated at 'Lys-14' by CREBBP (PubMed: [12789342](http://www.uniprot.org/citations/12789342)). Additionally, phosphorylates the CREBBP-interacting protein NCOA3. Also phosphorylates FOXO3 and may regulate this pro- apoptotic transcription factor (PubMed: [15084260](http://www.uniprot.org/citations/15084260)). Phosphorylates RIPK1 at 'Ser-25' which represses its kinase activity and consequently prevents TNF-mediated RIPK1-dependent cell death (By similarity). Phosphorylates AMBRA1 following mitophagy induction, promoting AMBRA1 interaction with ATG8 family proteins and its mitophagic activity (PubMed: [30217973](http://www.uniprot.org/citations/30217973)).

Cellular Location

Cytoplasm. Nucleus Note=Shuttles between the cytoplasm and the nucleus

Tissue Location

Widely expressed.

CHUK Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CHUK Antibody - Images

Figure 1: Western blot analysis using CHUK mouse mAb against Raji (1), Jurkat (2), THP-1 (3) and K562 (4) cell lysate.




Figure 2: Immunofluorescence analysis of NIH/3T3 cells using CHUK mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 3: Flow cytometric analysis of A549 cells using CHUK mouse mAb (blue) and negative control (red).

CHUK Antibody - References

1. Mol Cancer. 2010 Jan 5;9:1. 2. J Infect Dis. 2010 May 1;201(9):1371-80.