

**PDK1 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1506a**

**Specification**

---

**PDK1 Antibody - Product Information**

Application	WB, IHC, IF, FC
Primary Accession	<a href="#">Q15118</a>
Reactivity	Human, Rat, Monkey
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	44kDa KDa

**Description**

Pyruvate dehydrogenase (PDH) is a mitochondrial multienzyme complex that catalyzes the oxidative decarboxylation of pyruvate and is one of the major enzymes responsible for the regulation of homeostasis of carbohydrate fuels in mammals. The enzymatic activity is regulated by a phosphorylation/dephosphorylation cycle. Phosphorylation of PDH by a specific pyruvate dehydrogenase kinase (PDK) results in inactivation. (provided by RefSeq). Tissue specificity: Expressed predominantly in the heart.

**Immunogen**

Purified recombinant fragment of human PDK1 expressed in E. Coli.

**Formulation**

Ascitic fluid containing 0.03% sodium azide. <br />

**PDK1 Antibody - Additional Information**

**Gene ID** 5163

**Other Names**

[Pyruvate dehydrogenase (acetyl-transferring)] kinase isozyme 1, mitochondrial, 2.7.11.2, Pyruvate dehydrogenase kinase isoform 1, PDH kinase 1, PDK1, PDHK1

**Dilution**

WB~~1/500 - 1/2000  
IHC~~1/200 - 1/1000  
IF~~1/200 - 1/1000  
FC~~1/200 - 1/400

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PDK1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## PDK1 Antibody - Protein Information

**Name** PDK1

**Synonyms** PDHK1

### Function

Kinase that plays a key role in regulation of glucose and fatty acid metabolism and homeostasis via phosphorylation of the pyruvate dehydrogenase subunits PDHA1 and PDHA2. This inhibits pyruvate dehydrogenase activity, and thereby regulates metabolite flux through the tricarboxylic acid cycle, down-regulates aerobic respiration and inhibits the formation of acetyl-coenzyme A from pyruvate. Plays an important role in cellular responses to hypoxia and is important for cell proliferation under hypoxia. Protects cells against apoptosis in response to hypoxia and oxidative stress.

### Cellular Location

Mitochondrion matrix

### Tissue Location

Expressed predominantly in the heart. Detected at lower levels in liver, skeletal muscle and pancreas

## PDK1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## PDK1 Antibody - Images




Figure 1: Western blot analysis using PDK1 mouse mAb against NIH/3T3 (1), HeLa (2), Jurkat (3), HepG2 (4), PC-12 (5), and Cos7 (6) cell lysate.




Figure 2: Immunohistochemical analysis of paraffin-embedded breast cancer tissues (left) and brain tissues (right) using PDK1 mouse mAb with DAB staining.




Figure 3: Immunofluorescence analysis of HELA cells using PDK1 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 4: Flow cytometric analysis of Lovo cells using PDK1 mouse mAb (green) and negative control (purple).

#### **PDK1 Antibody - References**

1. Nat Cell Biol. 2008 Feb;10(2):127-37.
2. Blood. 2008 Apr 1;111(7):3723-34.
3. J Biol Chem. 2007 Apr 20;282(16):12272-89.