

OTX2 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1567a**Specification****OTX2 Antibody - Product Information**

Application	E, WB, IHC, IF, FC
Primary Accession	P32243
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	32kDa KDa

Description

This gene encodes a member of the bicoid sub-family of homeodomain-containing transcription factors. The encoded protein acts as a transcription factor and may play a role in brain and sensory organ development. A similar protein in mice is required for proper forebrain development. Tissue specificity: Expressed in brain.

Immunogen

Purified recombinant fragment of human OTX2 expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

OTX2 Antibody - Additional Information

Gene ID 5015

Other Names

Homeobox protein OTX2, Orthodenticle homolog 2, OTX2

Dilution

E~~1/10000
WB~~1/500 - 1/2000
IHC~~1/200 - 1/1000
IF~~1/200 - 1/1000
FC~~1/200 - 1/400

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OTX2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

OTX2 Antibody - Protein Information

Name OTX2

Function

Transcription factor probably involved in the development of the brain and the sense organs. Can bind to the bicoid/BCD target sequence (BTS): 5'-TCTAATCCC-3'.

Cellular Location

Nucleus.

OTX2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)





Figure 1: Western blot analysis using OTX2 mouse mAb against HepG2 (1), Jurkat (2), and NTERA-2 (3) cell lysate.




Figure 2: Western blot analysis using OTX2 mAb against human OTX2 (AA: 40-297) recombinant protein. (Expected MW is 65 kDa)




Figure 3: Immunohistochemical analysis of paraffin-embedded prostate tissues (left) and colon cancer tissues (right) using OTX2 mouse mAb with DAB staining.




Figure 4: Immunofluorescence analysis of HepG2 cells using OTX2 mouse mAb (green). Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 5: Flow cytometric analysis of HepG2 cells using OTX2 mouse mAb (green) and negative control (purple).

OTX2 Antibody - References

1. Hum Mutat. 2008 Nov;29(11):E278-83.
2. Cancer Res. 2010 Jan 1;70(1):181-91.