

**FOXO1 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1677a****Specification****FOXO1 Antibody - Product Information**

Application	<b>E, WB, IHC, IF</b>
Primary Accession	<a href="#">O12778</a>
Reactivity	<b>Human, Mouse</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG1</b>
Calculated MW	<b>69.7kDa KDa</b>

**Description**

This gene belongs to the forkhead family of transcription factors which are characterized by a distinct forkhead domain. The specific function of this gene has not yet been determined; however, it may play a role in myogenic growth and differentiation. Translocation of this gene with PAX3 has been associated with alveolar rhabdomyosarcoma.

**Immunogen**

Purified recombinant fragment of human FOXO1 expressed in E. Coli. <br />

**Formulation**

Purified antibody in PBS with 0.05% sodium azide

**FOXO1 Antibody - Additional Information**

**Gene ID** 2308

**Other Names**

Forkhead box protein O1, Forkhead box protein O1A, Forkhead in rhabdomyosarcoma, FOXO1, FKHR, FOXO1A

**Dilution**

E~~1/10000  
WB~~1/500 - 1/2000  
IHC~~1/200 - 1/1000  
IF~~1/200 - 1/1000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

FOXO1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**FOXO1 Antibody - Protein Information**

**Name** FOXO1 {ECO:0000303|PubMed:12228231, ECO:0000312|HGNC:HGNC:3819}

### Function

Transcription factor that is the main target of insulin signaling and regulates metabolic homeostasis in response to oxidative stress (PubMed:<a href="http://www.uniprot.org/citations/10358076" target="\_blank">10358076</a>, PubMed:<a href="http://www.uniprot.org/citations/12228231" target="\_blank">12228231</a>, PubMed:<a href="http://www.uniprot.org/citations/15220471" target="\_blank">15220471</a>, PubMed:<a href="http://www.uniprot.org/citations/15890677" target="\_blank">15890677</a>, PubMed:<a href="http://www.uniprot.org/citations/18356527" target="\_blank">18356527</a>, PubMed:<a href="http://www.uniprot.org/citations/19221179" target="\_blank">19221179</a>, PubMed:<a href="http://www.uniprot.org/citations/20543840" target="\_blank">20543840</a>, PubMed:<a href="http://www.uniprot.org/citations/21245099" target="\_blank">21245099</a>). Binds to the insulin response element (IRE) with consensus sequence 5'-TT[G/A]TTTTG-3' and the related Daf-16 family binding element (DBE) with consensus sequence 5'-TT[G/A]TTTAC-3' (PubMed:<a href="http://www.uniprot.org/citations/10358076" target="\_blank">10358076</a>). Activity suppressed by insulin (PubMed:<a href="http://www.uniprot.org/citations/10358076" target="\_blank">10358076</a>). Main regulator of redox balance and osteoblast numbers and controls bone mass (By similarity). Orchestrates the endocrine function of the skeleton in regulating glucose metabolism (By similarity). Also acts as a key regulator of chondrogenic commitment of skeletal progenitor cells in response to lipid availability: when lipids levels are low, translocates to the nucleus and promotes expression of SOX9, which induces chondrogenic commitment and suppresses fatty acid oxidation (By similarity). Acts synergistically with ATF4 to suppress osteocalcin/BGLAP activity, increasing glucose levels and triggering glucose intolerance and insulin insensitivity (By similarity). Also suppresses the transcriptional activity of RUNX2, an upstream activator of osteocalcin/BGLAP (By similarity). Acts as an inhibitor of glucose sensing in pancreatic beta cells by acting as a transcription repressor and suppressing expression of PDX1 (By similarity). In hepatocytes, promotes gluconeogenesis by acting together with PPARGC1A and CEBPA to activate the expression of genes such as IGFBP1, G6PC1 and PCK1 (By similarity). Also promotes gluconeogenesis by directly promoting expression of PPARGC1A and G6PC1 (PubMed:<a href="http://www.uniprot.org/citations/17024043" target="\_blank">17024043</a>). Important regulator of cell death acting downstream of CDK1, PKB/AKT1 and STK4/MST1 (PubMed:<a href="http://www.uniprot.org/citations/18356527" target="\_blank">18356527</a>, PubMed:<a href="http://www.uniprot.org/citations/19221179" target="\_blank">19221179</a>). Promotes neural cell death (PubMed:<a href="http://www.uniprot.org/citations/18356527" target="\_blank">18356527</a>). Mediates insulin action on adipose tissue (By similarity). Regulates the expression of adipogenic genes such as PPARG during preadipocyte differentiation and, adipocyte size and adipose tissue-specific gene expression in response to excessive calorie intake (By similarity). Regulates the transcriptional activity of GADD45A and repair of nitric oxide-damaged DNA in beta-cells (By similarity). Required for the autophagic cell death induction in response to starvation or oxidative stress in a transcription-independent manner (PubMed:<a href="http://www.uniprot.org/citations/20543840" target="\_blank">20543840</a>). Mediates the function of MLIP in cardiomyocytes hypertrophy and cardiac remodeling (By similarity). Positive regulator of apoptosis in cardiac smooth muscle cells as a result of its transcriptional activation of pro-apoptotic genes (PubMed:<a href="http://www.uniprot.org/citations/19483080" target="\_blank">19483080</a>). Regulates endothelial cell (EC) viability and apoptosis in a PPIA/CYPA- dependent manner via transcription of CCL2 and BCL2L1 which are involved in EC chemotaxis and apoptosis (PubMed:<a href="http://www.uniprot.org/citations/31063815" target="\_blank">31063815</a>).

### Cellular Location

Cytoplasm. Nucleus Note=Shuttles between the cytoplasm and nucleus. Largely nuclear in unstimulated cells (PubMed:11311120, PubMed:12228231, PubMed:19221179, PubMed:20543840, PubMed:21245099, PubMed:25009184). In osteoblasts, colocalizes with ATF4 and RUNX2 in the nucleus (By similarity). Serum deprivation increases localization to the nucleus, leading to activate expression of SOX9 and subsequent chondrogenesis (By similarity) Insulin-induced phosphorylation at Ser-256 by PKB/AKT1 leads, via stimulation of Thr-24 phosphorylation, to

binding of 14-3-3 proteins and nuclear export to the cytoplasm where it is degraded by the ubiquitin-proteasomal pathway (PubMed:11237865, PubMed:12228231) Phosphorylation at Ser-249 by CDK1 disrupts binding of 14-3-3 proteins and promotes nuclear accumulation (PubMed:18356527). Phosphorylation by NLK results in nuclear export (By similarity). Translocates to the nucleus upon oxidative stress-induced phosphorylation at Ser-212 by STK4/MST1 (PubMed:19221179, PubMed:21245099). SGK1-mediated phosphorylation also results in nuclear translocation (By similarity) Retained in the nucleus under stress stimuli including oxidative stress, nutrient deprivation or nitric oxide (By similarity). Retained in the nucleus on methylation (By similarity). PPIA/CYPA stimulates its nuclear accumulation (PubMed:31063815). Deacetylation by SIRT6, promotes its translocation into the cytoplasm (PubMed:25009184)

{ECO:0000250|UniProtKB:Q9R1E0, ECO:0000269|PubMed:11237865, ECO:0000269|PubMed:11311120, ECO:0000269|PubMed:12228231, ECO:0000269|PubMed:18356527, ECO:0000269|PubMed:19221179, ECO:0000269|PubMed:20543840, ECO:0000269|PubMed:21245099, ECO:0000269|PubMed:25009184, ECO:0000269|PubMed:31063815}


### Tissue Location

Expressed in umbilical endothelial cells (at protein level) (PubMed:19483080). Abundantly expressed in skeletal muscle and ovary, with lower expression in the heart, placenta, lung, liver, pancreas, spleen, testis and small intestine (PubMed:9479491) Weakly expressed in the brain, thymus, prostate and mucosal lining of the colon (PubMed:9479491).

### FOXO1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 2: Western blot analysis using FOXO1 mAb against human FOXO1 (AA: 471-600) recombinant protein. (Expected MW is 39.3 kDa)




Figure 3: Western blot analysis using FOXO1 mouse mAb against Hela (1), HEK293 (2), MCF-7(3), and C6 (4) cell lysate.




Figure 4: Immunohistochemical analysis of paraffin-embedded intima cancer tissues using FOXO1 mouse mAb with DAB staining.




Figure 5: Immunofluorescence analysis of HeLa cells using FOXO1 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye.

#### **FOXO1 Antibody - References**

Int J Oncol. 2009 Nov;35(5):1045-51 Cancer Res. 2009 Jul 1;69(13):5433-40.