

CK5 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1699a

Specification

CK5 Antibody - Product Information

Application	E, WB, IHC, IF, FC
Primary Accession	P13647
Reactivity	Human, Mouse, Monkey
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	62.3kDa KDa

Description

The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the basal layer of the epidermis with family member KRT14. Mutations in these genes have been associated with a complex of diseases termed epidermolysis bullosa simplex. The type II cytokeratins are clustered in a region of chromosome 12q12-q13.

Immunogen

Purified recombinant fragment of human CK5 expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

CK5 Antibody - Additional Information

Gene ID 3852

Other Names

Keratin, type II cytoskeletal 5, 58 kDa cytokeratin, Cytokeratin-5, CK-5, Keratin-5, K5, Type-II keratin Kb5, KRT5

Dilution

E~~1/10000
WB~~1/500 - 1/2000
IHC~~1/200 - 1/1000
IF~~1/200 - 1/1000
FC~~1/200 - 1/400

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CK5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CK5 Antibody - Protein Information

Name KRT5

Function

Required for the formation of keratin intermediate filaments in the basal epidermis and maintenance of the skin barrier in response to mechanical stress (By similarity). Regulates the recruitment of Langerhans cells to the epidermis, potentially by modulation of the abundance of macrophage chemotactic cytokines, macrophage inflammatory cytokines and CTNND1 localization in keratinocytes (By similarity).

Cellular Location

Cytoplasm.


Tissue Location

Expressed in corneal epithelium (at protein level) (PubMed:26758872). Expressed in keratinocytes (at protein level) (PubMed:20128788, PubMed:31302245).

CK5 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using CK5 mAb against human CK5 (AA: 158-272) recombinant protein. (Expected MW is 33.3 kDa)




Figure 2: Western blot analysis using CK5 mouse mAb against A431 (1), MCF-7 (2), HeLa (3), HepG2 (4), 3T3-L1 (5), and COS-7 (6) cell lysate.




Figure 3: Immunohistochemical analysis of paraffin-embedded breast cancer tissues using CK5 mouse mAb with DAB staining.




Figure 4: Immunohistochemical analysis of paraffin-embedded stomach cancer tissues using CK5 mouse mAb with DAB staining.




Figure 5: Immunofluorescence analysis of HeLa cells. Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 6: Immunofluorescence analysis of HeLa cells using CK5 mouse mAb (green). Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 7: Flow cytometric analysis of HeLa cells using CK5 mouse mAb (green) and negative control (red).

CK5 Antibody - References

Am J Surg Pathol. 2009 Nov;33(11):1615-23. J Dermatol. 2009 Aug;36(8):447-52.