

**MAP1LC3B Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1713a**

**Specification**

**MAP1LC3B Antibody - Product Information**

Application	<b>E, WB, FC</b>
Primary Accession	<a href="#">O9GZQ8</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG1</b>
Calculated MW	<b>14.6kDa KDa</b>

**Description**

The product of this gene is a subunit of neuronal microtubule-associated MAP1A and MAP1B proteins, which are involved in microtubule assembly and important for neurogenesis. Studies on the rat homolog implicate a role for this gene in autophagy, a process that involves the bulk degradation of cytoplasmic component.

**Immunogen**

Purified recombinant fragment of human MAP1LC3B expressed in E. Coli. <br />

**Formulation**

Ascitic fluid containing 0.03% sodium azide. <br />

**MAP1LC3B Antibody - Additional Information**

**Gene ID** 81631

**Other Names**

Microtubule-associated proteins 1A/1B light chain 3B, Autophagy-related protein LC3 B, Autophagy-related ubiquitin-like modifier LC3 B, MAP1 light chain 3-like protein 2, MAP1A/MAP1B light chain 3 B, MAP1A/MAP1B LC3 B, Microtubule-associated protein 1 light chain 3 beta, MAP1LC3B, MAP1ALC3

**Dilution**

E~~1/10000  
WB~~1/500 - 1/2000  
FC~~1/200 - 1/400

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

MAP1LC3B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## MAP1LC3B Antibody - Protein Information

Name MAP1LC3B ([HGNC:13352](#))

Synonyms MAP1ALC3

### Function

Ubiquitin-like modifier involved in formation of autophagosomal vacuoles (autophagosomes) (PubMed:[20418806](http://www.uniprot.org/citations/20418806)), PubMed:[23209295](http://www.uniprot.org/citations/23209295), PubMed:[28017329](http://www.uniprot.org/citations/28017329)). Plays a role in mitophagy which contributes to regulate mitochondrial quantity and quality by eliminating the mitochondria to a basal level to fulfill cellular energy requirements and preventing excess ROS production (PubMed:[23209295](http://www.uniprot.org/citations/23209295), PubMed:[28017329](http://www.uniprot.org/citations/28017329)). In response to cellular stress and upon mitochondria fission, binds C-18 ceramides and anchors autophagolysosomes to outer mitochondrial membranes to eliminate damaged mitochondria (PubMed:[22922758](http://www.uniprot.org/citations/22922758)). While LC3s are involved in elongation of the phagophore membrane, the GABARAP/GATE-16 subfamily is essential for a later stage in autophagosome maturation (PubMed:[20418806](http://www.uniprot.org/citations/20418806), PubMed:[23209295](http://www.uniprot.org/citations/23209295), PubMed:[28017329](http://www.uniprot.org/citations/28017329)). Promotes primary ciliogenesis by removing OFD1 from centriolar satellites via the autophagic pathway (PubMed:[24089205](http://www.uniprot.org/citations/24089205)). Through its interaction with the reticulophagy receptor TEX264, participates in the remodeling of subdomains of the endoplasmic reticulum into autophagosomes upon nutrient stress, which then fuse with lysosomes for endoplasmic reticulum turnover (PubMed:[31006537](http://www.uniprot.org/citations/31006537), PubMed:[31006538](http://www.uniprot.org/citations/31006538)). Upon nutrient stress, directly recruits cofactor JMY to the phagophore membrane surfaces and promotes JMY's actin nucleation activity and autophagosome biogenesis during autophagy (PubMed:[30420355](http://www.uniprot.org/citations/30420355)).

### Cellular Location

Cytoplasmic vesicle, autophagosome membrane; Lipid-anchor Endomembrane system; Lipid-anchor Mitochondrion membrane; Lipid-anchor. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:Q9CQV6}. Cytoplasmic vesicle. Note=LC3-II binds to the autophagic membranes. LC3-II localizes with the mitochondrial inner membrane during Parkin-mediated mitophagy (PubMed:28017329). Localizes also to discrete punctae along the ciliary axoneme

### Tissue Location

Most abundant in heart, brain, skeletal muscle and testis. Little expression observed in liver

## MAP1LC3B Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)





Figure 1: Western blot analysis using MAP1LC3B mAb against human MAP1LC3B (AA: 1-125) recombinant protein. (Expected MW is 40.2 kDa)




Figure 2: Western blot analysis using MAP1LC3B mAb against HEK293 (1) and MAP1LC3B (AA: 1-125)-hlgGfc transfected HEK293 (2) cell lysate.




Figure 3: Flow cytometric analysis of HeLa cells using MAP1LC3B mouse mAb (green) and negative control (red).

### MAP1LC3B Antibody - References

1. J Clin Lab Anal. 2009;23(4):249-58.
2. Ai Zheng. 2008 Jan;27(1):25-9.