

FZD5 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1760a**Specification****FZD5 Antibody - Product Information**

Application	E, WB, FC
Primary Accession	Q13467
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	64.5kDa KDa

Description

Members of the 'frizzled' gene family encode 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The FZD5 protein is believed to be the receptor for the Wnt5A ligand.

Immunogen

Purified recombinant fragment of human FZD5 (AA:151-217) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

FZD5 Antibody - Additional Information

Gene ID 7855

Other Names

Frizzled-5, Fz-5, hFz5, FzE5, FZD5, C2orf31

Dilution

E~~1/10000
WB~~1/500 - 1/2000
FC~~1/200 - 1/400

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FZD5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

FZD5 Antibody - Protein Information

Name FZD5

Synonyms C2orf31

Function

Receptor for Wnt proteins (PubMed: [10097073](http://www.uniprot.org/citations/10097073), PubMed: [20530549](http://www.uniprot.org/citations/20530549) target="_blank">20530549, PubMed: [26908622](http://www.uniprot.org/citations/26908622) target="_blank">26908622, PubMed: [9054360](http://www.uniprot.org/citations/9054360) target="_blank">9054360). Functions in the canonical Wnt/beta- catenin signaling pathway. In vitro activates WNT2, WNT10B, WNT5A, but not WNT2B or WNT4 signaling (By similarity). In neurons, activation by WNT7A promotes formation of synapses (PubMed: [20530549](http://www.uniprot.org/citations/20530549)). May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues (Probable). Plays a role in yolk sac angiogenesis and in placental vascularization (By similarity). Plays a role in ocular development (PubMed: [26908622](http://www.uniprot.org/citations/26908622)).


Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q8CHL0}. Golgi apparatus membrane {ECO:0000250|UniProtKB:Q9EQD0}; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q9EQD0}. Synapse {ECO:0000250|UniProtKB:Q8CHL0}. Perikaryon {ECO:0000250|UniProtKB:Q8CHL0}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q8CHL0}. Cell projection, axon {ECO:0000250|UniProtKB:Q8CHL0}. Note=Localized at the plasma membrane and also found at the Golgi apparatus. {ECO:0000250|UniProtKB:Q9EQD0}

FZD5 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using FZD5 mAb against human FZD5 recombinant protein. (Expected MW is 32.5 kDa)




Figure 2: Western blot analysis using FZD5 mouse mAb against A549 (1), and PC-3 (2) cell lysate.




Figure 3: Flow cytometric analysis of HepG2 cells using FZD5 mouse mAb (green) and negative control (red).

FZD5 Antibody - References

1. J Biol Chem. 2009 Sep 25;284(39):26716-24. 2. Int J Oncol. 2007 Mar;30(3):751-5.