

**CD22 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1851a****Specification****CD22 Antibody - Product Information**

Application	<b>E, WB, IF, FC</b>
Primary Accession	<a href="#">P20273</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG1</b>
Calculated MW	<b>95.3kDa KDa</b>

**Description**

CD22 may be involved in the localization of B-cells in lymphoid tissues. Binds sialylated glycoproteins; one of which is CD45. Preferentially binds to alpha-2,6-linked sialic acid. The sialic acid recognition site can be masked by cis interactions with sialic acids on the same cell surface. Upon ligand induced tyrosine phosphorylation in the immune response seems to be involved in regulation of B-cell antigen receptor signaling. Plays a role in positive regulation through interaction with Src family tyrosine kinases and may also act as an inhibitory receptor by recruiting cytoplasmic phosphatases via their SH2 domains that block signal transduction through dephosphorylation of signaling molecules

**Immunogen**

Purified recombinant fragment of human CD22 (AA: 621-725) expressed in E. Coli.

**Formulation**

Purified antibody in PBS with 0.05% sodium azide

**CD22 Antibody - Additional Information**

**Gene ID** 933

**Other Names**

B-cell receptor CD22, B-lymphocyte cell adhesion molecule, BL-CAM, Sialic acid-binding Ig-like lectin 2, Siglec-2, T-cell surface antigen Leu-14, CD22, CD22, SIGLEC2

**Dilution**

E~~1/10000  
WB~~1/500 - 1/2000  
IF~~1/200 - 1/1000  
FC~~1/200 - 1/400

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CD22 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## CD22 Antibody - Protein Information

**Name** CD22 {ECO:0000303|PubMed:1691828, ECO:0000312|HGNC:HGNC:1643}

### Function

Most highly expressed siglec (sialic acid-binding immunoglobulin-like lectin) on B-cells that plays a role in various aspects of B-cell biology including differentiation, antigen presentation, and trafficking to bone marrow (PubMed:<a href="http://www.uniprot.org/citations/8627166" target="\_blank">8627166</a>, PubMed:<a href="http://www.uniprot.org/citations/34330755" target="\_blank">34330755</a>). Binds to alpha 2,6-linked sialic acid residues of surface molecules such as CD22 itself, CD45 and IgM in a cis configuration. Can also bind to ligands on other cells as an adhesion molecule in a trans configuration (PubMed:<a href="http://www.uniprot.org/citations/20172905" target="\_blank">20172905</a>). Acts as an inhibitory coreceptor on the surface of B-cells and inhibits B-cell receptor induced signaling, characterized by inhibition of the calcium mobilization and cellular activation. Mechanistically, the immunoreceptor tyrosine-based inhibitory motif domain is phosphorylated by the Src kinase LYN, which in turn leads to the recruitment of the protein tyrosine phosphatase 1/PTPN6, leading to the negative regulation of BCR signaling (PubMed:<a href="http://www.uniprot.org/citations/8627166" target="\_blank">8627166</a>). If this negative signaling from is of sufficient strength, apoptosis of the B-cell can be induced (PubMed:<a href="http://www.uniprot.org/citations/20516366" target="\_blank">20516366</a>).

### Cellular Location

Cell membrane; Single-pass type I membrane protein

### Tissue Location

B-lymphocytes.

## CD22 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)





Figure 1: Western blot analysis using CD22 mAb against human CD22 recombinant protein. (Expected MW is 37 kDa)




Figure 2: Western blot analysis using CD22 mAb against HEK293 (1) and CD22 (AA: 621-725)-hIgGfc transfected HEK293 (2) cell lysate.




Figure 3: Immunofluorescence analysis of HeLa cells using CD22 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 4: Flow cytometric analysis of HeLa cells using CD22 mouse mAb (green) and negative control (red).

### CD22 Antibody - Background

CD22 may be involved in the localization of B-cells in lymphoid tissues. Binds sialylated glycoproteins; one of which is CD45. Preferentially binds to alpha-2,6-linked sialic acid. The sialic acid recognition site can be masked by cis interactions with sialic acids on the same cell surface. Upon ligand induced tyrosine phosphorylation in the immune response seems to be involved in regulation of B-cell antigen receptor signaling. Plays a role in positive regulation through interaction with Src family tyrosine kinases and may also act as an inhibitory receptor by recruiting cytoplasmic phosphatases via their SH2 domains that block signal transduction through dephosphorylation of signaling molecules ;

### CD22 Antibody - References

1. Cancer Res. 2012 Nov 1;72(21):5556-65.
2. J Innate Immun. 2011;3(4):411-9.