

**CD14 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1886a**

**Specification**

**CD14 Antibody - Product Information**

Application	<b>E, WB, IF, IHC</b>
Primary Accession	<a href="#">P08571</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG2a</b>
Calculated MW	<b>40kDa KDa</b>

**Description**

The protein encoded by this gene is a surface antigen that is preferentially expressed on monocytes/macrophages. It cooperates with other proteins to mediate the innate immune response to bacterial lipopolysaccharide. Alternative splicing results in multiple transcript variants encoding the same protein.

**Immunogen**

Purified recombinant fragment of human CD14 (AA: 20-214) expressed in E. Coli.

**Formulation**

Purified antibody in PBS with 0.05% sodium azide

**CD14 Antibody - Additional Information**

**Gene ID** 929

**Other Names**

Monocyte differentiation antigen CD14, Myeloid cell-specific leucine-rich glycoprotein, CD14, Monocyte differentiation antigen CD14, urinary form, Monocyte differentiation antigen CD14, membrane-bound form, CD14

**Dilution**

E~~1/10000  
WB~~1/500 - 1/2000  
IF~~1/200 - 1/1000  
IHC~~1/200 - 1/1000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CD14 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**CD14 Antibody - Protein Information**

**Name** CD14**Function**

Coreceptor for bacterial lipopolysaccharide (PubMed:<a href="http://www.uniprot.org/citations/1698311" target="\_blank">1698311</a>, PubMed:<a href="http://www.uniprot.org/citations/23264655" target="\_blank">23264655</a>). In concert with LBP, binds to monomeric lipopolysaccharide and delivers it to the LY96/TLR4 complex, thereby mediating the innate immune response to bacterial lipopolysaccharide (LPS) (PubMed:<a href="http://www.uniprot.org/citations/20133493" target="\_blank">20133493</a>, PubMed:<a href="http://www.uniprot.org/citations/22265692" target="\_blank">22265692</a>, PubMed:<a href="http://www.uniprot.org/citations/23264655" target="\_blank">23264655</a>). Acts via MyD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed:<a href="http://www.uniprot.org/citations/8612135" target="\_blank">8612135</a>). Acts as a coreceptor for TLR2:TLR6 heterodimer in response to diacylated lipopeptides and for TLR2:TLR1 heterodimer in response to triacylated lipopeptides, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway (PubMed:<a href="http://www.uniprot.org/citations/16880211" target="\_blank">16880211</a>). Binds electronegative LDL (LDL(-)) and mediates the cytokine release induced by LDL(-) (PubMed:<a href="http://www.uniprot.org/citations/23880187" target="\_blank">23880187</a>).

**Cellular Location**

Cell membrane; Lipid-anchor, GPI-anchor. Secreted. Membrane raft. Golgi apparatus.  
Note=Secreted forms may arise by cleavage of the GPI anchor.

**Tissue Location**

Detected on macrophages (at protein level) (PubMed:1698311). Expressed strongly on the surface of monocytes and weakly on the surface of granulocytes; also expressed by most tissue macrophages.

**CD14 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)





Figure 1: Western blot analysis using CD14 mAb against human CD14 (AA: 20-214) recombinant protein. (Expected MW is 46.8 kDa)




Figure 2: Western blot analysis using CD14 mAb against HEK293 (1) and CD14 (AA: 20-214)-hlgGfc transfected HEK293 (2) cell lysate.




Figure 3: Immunofluorescence analysis of HepG2 cells using CD14 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Secondary antibody from Fisher (Cat#: 35503)




Figure 4: Immunohistochemical analysis of paraffin-embedded cervical cancer tissues using CD14 mouse mAb with DAB staining.




Figure 5: Immunohistochemical analysis of paraffin-embedded colon tissues using CD14 mouse mAb with DAB staining.

### CD14 Antibody - Background

The membrane-associated protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MDR/TAP subfamily. Members of the MDR/TAP subfamily are involved in multidrug resistance. The protein encoded by this gene is an

ATP-dependent drug efflux pump for xenobiotic compounds with broad substrate specificity. It is responsible for decreased drug accumulation in multidrug-resistant cells and often mediates the development of resistance to anticancer drugs. This protein also functions as a transporter in the blood-brain barrier. ;

#### **CD14 Antibody - References**

1. J Immunol. 2012 Dec 15;189(12):5729-44.
2. Iran J Immunol. 2011 Jun;8(2):111-9.