

CAMK2G Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1916a

Specification

CAMK2G Antibody - Product Information

Application	E, WB
Primary Accession	O13555
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	62.6kDa KDa

Description

The product of this gene is one of the four subunits of an enzyme which belongs to the serine/threonine protein kinase family, and to the Ca(2+)/calmodulin-dependent protein kinase subfamily. Calcium signaling is crucial for several aspects of plasticity at glutamatergic synapses. In mammalian cells the enzyme is composed of four different chains: alpha, beta, gamma, and delta. The product of this gene is a gamma chain. Many alternatively spliced transcripts encoding different isoforms have been described but the full-length nature of all the variants has not been determined.

Immunogen

Purified recombinant fragment of human CAMK2G (AA: 322-481) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide.

CAMK2G Antibody - Additional Information

Gene ID 818

Other Names

Calcium/calmodulin-dependent protein kinase type II subunit gamma, CaM kinase II subunit gamma, CaMK-II subunit gamma, 2.7.11.17, CAMK2G, CAMK, CAMK-II, CAMKG

Dilution

E~~1/10000
WB~~1/500 - 1/2000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CAMK2G Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CAMK2G Antibody - Protein Information

Name CAMK2G**Synonyms** CAMK, CAMK-II, CAMKG**Function**

Calcium/calmodulin-dependent protein kinase that functions autonomously after Ca(2+)/calmodulin-binding and autophosphorylation, and is involved in sarcoplasmic reticulum Ca(2+) transport in skeletal muscle and may function in dendritic spine and synapse formation and neuronal plasticity (PubMed:16690701). In slow-twitch muscles, is involved in regulation of sarcoplasmic reticulum (SR) Ca(2+) transport and in fast-twitch muscle participates in the control of Ca(2+) release from the SR through phosphorylation of the ryanodine receptor-coupling factor triadin (PubMed:16690701). In the central nervous system, it is involved in the regulation of neurite formation and arborization (PubMed:30184290). It may participate in the promotion of dendritic spine and synapse formation and maintenance of synaptic plasticity which enables long-term potentiation (LTP) and hippocampus-dependent learning. In response to interferon-gamma (IFN-gamma) stimulation, catalyzes phosphorylation of STAT1, stimulating the JAK-STAT signaling pathway (By similarity).

Cellular Location

Sarcoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side


Tissue Location

Expressed in skeletal muscle.

CAMK2G Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using CAMK2G mAb against human CAMK2G (AA: 322-481) recombinant protein. (Expected MW is 44 kDa)




Figure 2: Western blot analysis using CAMK2G mAb against HEK293 (1) and CAMK2G (AA: 322-481)-hlgGfc transfected HEK293 (2) cell lysate.

CAMK2G Antibody - Background

This gene encodes a member of the intermediate filament protein family and is expressed primarily in nerve cells. ; ;

CAMK2G Antibody - References

1. Blood. 2012 Dec 6;120(24):4829-39.
2. Diabetologia. 2002 Apr;45(4):580-3.