

**NOX4 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO2017a****Specification****NOX4 Antibody - Product Information**

Application	<b>E, WB, IF, FC, IHC</b>
Primary Accession	<a href="#">O9NPH5</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG1</b>
Calculated MW	<b>67kDa KDa</b>

**Description**

This gene encodes a member of the NOX-family of enzymes that functions as the catalytic subunit the NADPH oxidase complex. The encoded protein is localized to non-phagocytic cells where it acts as an oxygen sensor and catalyzes the reduction of molecular oxygen to various reactive oxygen species (ROS). The ROS generated by this protein have been implicated in numerous biological functions including signal transduction, cell differentiation and tumor cell growth. A pseudogene has been identified on the other arm of chromosome 11. Alternative splicing results in multiple transcript variants.

**Immunogen**

Purified recombinant fragment of human NOX4 (AA: 210-310) expressed in E. Coli.

**Formulation**

Purified antibody in PBS with 0.05% sodium azide

**NOX4 Antibody - Additional Information**

**Gene ID** 50507

**Other Names**

NADPH oxidase 4, 1.6.3.-, Kidney oxidase-1, KOX-1, Kidney superoxide-producing NADPH oxidase, Renal NAD(P)H-oxidase, NOX4, RENOX

**Dilution**

E~~1/10000  
WB~~1/500 - 1/2000  
IF~~1/200 - 1/1000  
FC~~1/200 - 1/400  
IHC~~1/200 - 1/1000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

NOX4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## NOX4 Antibody - Protein Information

**Name** NOX4

**Synonyms** RENOX

### Function

NADPH oxidase that catalyzes predominantly the reduction of oxygen to H<sub>2</sub>O<sub>2</sub> (PubMed:<a href="http://www.uniprot.org/citations/14966267" target="\_blank">14966267</a>, PubMed:<a href="http://www.uniprot.org/citations/15356101" target="\_blank">15356101</a>, PubMed:<a href="http://www.uniprot.org/citations/15927447" target="\_blank">15927447</a>, PubMed:<a href="http://www.uniprot.org/citations/21343298" target="\_blank">21343298</a>, PubMed:<a href="http://www.uniprot.org/citations/25062272" target="\_blank">25062272</a>). Can also catalyze to a smaller extent, the reduction of oxygen to superoxide (PubMed:<a href="http://www.uniprot.org/citations/10869423" target="\_blank">10869423</a>, PubMed:<a href="http://www.uniprot.org/citations/11032835" target="\_blank">11032835</a>, PubMed:<a href="http://www.uniprot.org/citations/15155719" target="\_blank">15155719</a>, PubMed:<a href="http://www.uniprot.org/citations/15572675" target="\_blank">15572675</a>, PubMed:<a href="http://www.uniprot.org/citations/15927447" target="\_blank">15927447</a>, PubMed:<a href="http://www.uniprot.org/citations/16019190" target="\_blank">16019190</a>, PubMed:<a href="http://www.uniprot.org/citations/16179589" target="\_blank">16179589</a>, PubMed:<a href="http://www.uniprot.org/citations/16230378" target="\_blank">16230378</a>, PubMed:<a href="http://www.uniprot.org/citations/16324151" target="\_blank">16324151</a>, PubMed:<a href="http://www.uniprot.org/citations/25062272" target="\_blank">25062272</a>). May function as an oxygen sensor regulating the KCNK3/TASK-1 potassium channel and HIF1A activity (PubMed:<a href="http://www.uniprot.org/citations/16019190" target="\_blank">16019190</a>). May regulate insulin signaling cascade (PubMed:<a href="http://www.uniprot.org/citations/14966267" target="\_blank">14966267</a>). May play a role in apoptosis, bone resorption and lipopolysaccharide-mediated activation of NFκB (PubMed:<a href="http://www.uniprot.org/citations/15356101" target="\_blank">15356101</a>, PubMed:<a href="http://www.uniprot.org/citations/15572675" target="\_blank">15572675</a>). May produce superoxide in the nucleus and play a role in regulating gene expression upon cell stimulation (PubMed:<a href="http://www.uniprot.org/citations/16324151" target="\_blank">16324151</a>).

### Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Cell junction, focal adhesion {ECO:0000250|UniProtKB:Q924V1}. Nucleus [Isoform 3]: Cytoplasm. Cytoplasm, perinuclear region [Isoform 6]: Cytoplasm. Cytoplasm, perinuclear region

### Tissue Location

Expressed by distal tubular cells in kidney cortex and in endothelial cells (at protein level). Widely expressed. Strongly expressed in kidney and to a lower extent in heart, adipocytes, hepatoma, endothelial cells, skeletal muscle, brain, several brain tumor cell lines and airway epithelial cells

## NOX4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)