

RALA Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO2091a

Specification

RALA Antibody - Product Information

Application	E, WB, FC
Primary Accession	P11233
Reactivity	Human, Mouse, Monkey
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	23.6kDa KDa

Description

The product of this gene belongs to the small GTPase superfamily, Ras family of proteins. GTP-binding proteins mediate the transmembrane signaling initiated by the occupancy of certain cell surface receptors. This gene encodes a low molecular mass ras-like GTP-binding protein that shares about 50% similarity with other ras proteins.

Immunogen

Purified recombinant fragment of human RALA (AA: 71-203) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

RALA Antibody - Additional Information

Gene ID 5898

Other Names

Ras-related protein Ral-A, RALA, RAL

Dilution

E~~1/10000
WB~~1/500 - 1/2000
FC~~1/200 - 1/400

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RALA Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

RALA Antibody - Protein Information

Name RALA

Synonyms RAL

Function

Multifunctional GTPase involved in a variety of cellular processes including gene expression, cell migration, cell proliferation, oncogenic transformation and membrane trafficking. Accomplishes its multiple functions by interacting with distinct downstream effectors (PubMed:18756269, PubMed:19306925, PubMed:20005108, PubMed:21822277, PubMed:30500825). Acts as a GTP sensor for GTP-dependent exocytosis of dense core vesicles. The RALA- exocyst complex regulates integrin-dependent membrane raft exocytosis and growth signaling (PubMed:20005108). Key regulator of LPAR1 signaling and competes with GRK2 for binding to LPAR1 thus affecting the signaling properties of the receptor. Required for anchorage- independent proliferation of transformed cells (PubMed:19306925). During mitosis, supports the stabilization and elongation of the intracellular bridge between dividing cells. Cooperates with EXOC2 to recruit other components of the exocyst to the early midbody (PubMed:18756269). During mitosis, also controls mitochondrial fission by recruiting to the mitochondrion RALBP1, which mediates the phosphorylation and activation of DNMI1 by the mitotic kinase cyclin B- CDK1 (PubMed:21822277).

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Cleavage furrow. Midbody, Midbody ring. Mitochondrion. Note=Predominantly at the cell surface in the absence of LPA. In the presence of LPA, colocalizes with LPAR1 and LPAR2 in endocytic vesicles (PubMed:19306925). May colocalize with CNTRL/centriolin at the midbody ring (PubMed:16213214). However, localization at the midbody at late cytokinesis was not confirmed (PubMed:18756269). Relocalizes to the mitochondrion during mitosis where it regulates mitochondrial fission (PubMed:21822277)

RALA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)