

**LEF1 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO2190a**

**Specification**

**LEF1 Antibody - Product Information**

Application	<b>E, WB, IHC</b>
Primary Accession	<a href="#">O9UJU2</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG1</b>
Calculated MW	<b>44.2kDa KDa</b>

**Description**

This gene encodes a transcription factor belonging to a family of proteins that share homology with the high mobility group protein-1. The protein encoded by this gene can bind to a functionally important site in the T-cell receptor-alpha enhancer, thereby conferring maximal enhancer activity. This transcription factor is involved in the Wnt signaling pathway, and it may function in hair cell differentiation and follicle morphogenesis. Mutations in this gene have been found in somatic sebaceous tumors. This gene has also been linked to other cancers, including androgen-independent prostate cancer. Alternative splicing results in multiple transcript variants.

**Immunogen**

Purified recombinant fragment of human LEF1 (AA: 33-138) expressed in E. Coli.

**Formulation**

Purified antibody in PBS with 0.05% sodium azide

**LEF1 Antibody - Additional Information**

**Gene ID** 51176

**Other Names**

Lymphoid enhancer-binding factor 1, LEF-1, T cell-specific transcription factor 1-alpha, TCF1-alpha, LEF1

**Dilution**

E~~1/10000  
WB~~1/500 - 1/2000  
IHC~~1/200 - 1/1000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

LEF1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## LEF1 Antibody - Protein Information

Name LEF1 ([HGNC:6551](#))

### Function

Transcription factor that binds DNA in a sequence-specific manner (PubMed:<a href="http://www.uniprot.org/citations/2010090" target="\_blank">2010090</a>). Participates in the Wnt signaling pathway (By similarity). Activates transcription of target genes in the presence of CTNNB1 and EP300 (By similarity). PIAG antagonizes both Wnt-dependent and Wnt-independent activation by LEF1 (By similarity). TLE1, TLE2, TLE3 and TLE4 repress transactivation mediated by LEF1 and CTNNB1 (PubMed:<a href="http://www.uniprot.org/citations/11266540" target="\_blank">11266540</a>). Regulates T-cell receptor alpha enhancer function (PubMed:<a href="http://www.uniprot.org/citations/19653274" target="\_blank">19653274</a>). Required for IL17A expressing gamma-delta T-cell maturation and development, via binding to regulator loci of BLK to modulate expression (By similarity). Acts as a positive regulator of odontoblast differentiation during mesenchymal tooth germ formation, expression is repressed during the bell stage by MSX1-mediated inhibition of CTNNB1 signaling (By similarity). May play a role in hair cell differentiation and follicle morphogenesis (By similarity).

### Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00267}. Note=Found in nuclear bodies upon PIASG binding.

### Tissue Location

Detected in thymus. Not detected in normal colon, but highly expressed in colon cancer biopsies and colon cancer cell lines. Expressed in several pancreatic tumors and weakly expressed in normal pancreatic tissue. Isoforms 1 and 5 are detected in several pancreatic cell lines.

## LEF1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)