

PCNA Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO2261a

Specification

PCNA Antibody - Product Information

Application	E, WB, FC, IHC
Primary Accession	P12004
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	28.7kDa KDa

Description

The protein encoded by this gene is found in the nucleus and is a cofactor of DNA polymerase delta. The encoded protein acts as a homotrimer and helps increase the processivity of leading strand synthesis during DNA replication. In response to DNA damage, this protein is ubiquitinated and is involved in the RAD6-dependent DNA repair pathway. Two transcript variants encoding the same protein have been found for this gene. Pseudogenes of this gene have been described on chromosome 4 and on the X chromosome.

Immunogen

Purified recombinant fragment of human PCNA (AA: 53-196) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

PCNA Antibody - Additional Information

Gene ID 5111

Other Names

Proliferating cell nuclear antigen, PCNA, Cyclin, PCNA

Dilution

E~~1/10000
WB~~1/500 - 1/2000
FC~~1/200 - 1/400
IHC~~1/200 - 1/1000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PCNA Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PCNA Antibody - Protein Information

Name PCNA

Function

Auxiliary protein of DNA polymerase delta and epsilon, is involved in the control of eukaryotic DNA replication by increasing the polymerase's processibility during elongation of the leading strand (PubMed:35585232). Induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase, but not apurinic-apyrimidinic (AP) endonuclease, APEX2 activities. Has to be loaded onto DNA in order to be able to stimulate APEX2. Plays a key role in DNA damage response (DDR) by being conveniently positioned at the replication fork to coordinate DNA replication with DNA repair and DNA damage tolerance pathways (PubMed:24939902). Acts as a loading platform to recruit DDR proteins that allow completion of DNA replication after DNA damage and promote postreplication repair: Monoubiquitinated PCNA leads to recruitment of translesion (TLS) polymerases, while 'Lys-63'-linked polyubiquitination of PCNA is involved in error-free pathway and employs recombination mechanisms to synthesize across the lesion (PubMed:24695737).


Cellular Location

Nucleus Note=Colocalizes with CREBBP, EP300 and POLD1 to sites of DNA damage (PubMed:24939902). Forms nuclear foci representing sites of ongoing DNA replication and vary in morphology and number during S phase (PubMed:15543136). Co-localizes with SMARCA5/SNF2H and BAZ1B/WSTF at replication foci during S phase (PubMed:15543136). Together with APEX2, is redistributed in discrete nuclear foci in presence of oxidative DNA damaging agents.

PCNA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using PCNA mAb against human PCNA recombinant protein. (Expected MW is 41.2 kDa)




Figure 2: Western blot analysis using PCNA mAb against HEK293 (1) and PCNA (AA: 53-196)-hIgGFc transfected HEK293 (2) cell lysate.




Figure 3: Western blot analysis using PCNA mouse mAb against A431 (1), HeLa (2), HepG2 (3), Raji (4), and MOLT4 (5) cell lysate.




Figure 4: Flow cytometric analysis of MOLT4 cells using PCNA mouse mAb (green) and negative control (purple).




Figure 5: Immunohistochemical analysis of paraffin-embedded cervical cancer tissues using PCNA mouse mAb with DAB staining.




Figure 6: Immunohistochemical analysis of paraffin-embedded colon cancer tissues using PCNA mouse mAb with DAB staining.

PCNA Antibody - References

- 1.Cancer Res. 2012 Jul 1;72(13):3217-27.
- 2.PLoS One. 2012;7(1):e29416.