

**Mouse Monoclonal Antibody to PYCARD**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO2344a**

**Specification**

---

**Mouse Monoclonal Antibody to PYCARD - Product Information**

Application	E, WB, FC, ICC, IHC
Primary Accession	<a href="#">O9ULZ3</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1
Calculated MW	21.6kDa KDa

**Description**

This gene encodes an adaptor protein that is composed of two protein-protein interaction domains: a N-terminal PYRIN-PAAD-DAPIN domain (PYD) and a C-terminal caspase-recruitment domain (CARD). The PYD and CARD domains are members of the six-helix bundle death domain-fold superfamily that mediates assembly of large signaling complexes in the inflammatory and apoptotic signaling pathways via the activation of caspase. In normal cells, this protein is localized to the cytoplasm; however, in cells undergoing apoptosis, it forms ball-like aggregates near the nuclear periphery. Two transcript variants encoding different isoforms have been found for this gene.;

**Immunogen**

Purified recombinant fragment of human PYCARD (AA: 1-120) expressed in E. Coli.

**Formulation**

Purified antibody in PBS with 0.05% sodium azide

**Application Note**

ELISA: 1/10000; WB: 1/500 - 1/2000; IHC: 1/200 - 1/1000; ICC: 1/200 - 1/1000; FCM: 1/200 - 1/400

**Mouse Monoclonal Antibody to PYCARD - Additional Information**

**Gene ID** 29108

**Other Names**

ASC; TMS; TMS1; CARD5; TMS-1

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Mouse Monoclonal Antibody to PYCARD is for research use only and not for use in diagnostic or therapeutic procedures.

**Mouse Monoclonal Antibody to PYCARD - Protein Information**

**Name** PYCARD {ECO:0000303|Ref.4, ECO:0000312|HGNC:HGNC:16608}

### Function

Functions as a key mediator in apoptosis and inflammation (PubMed:<a href="http://www.uniprot.org/citations/11103777" target="\_blank">11103777</a>, PubMed:<a href="http://www.uniprot.org/citations/12646168" target="\_blank">12646168</a>, PubMed:<a href="http://www.uniprot.org/citations/15030775" target="\_blank">15030775</a>, PubMed:<a href="http://www.uniprot.org/citations/17349957" target="\_blank">17349957</a>, PubMed:<a href="http://www.uniprot.org/citations/17599095" target="\_blank">17599095</a>, PubMed:<a href="http://www.uniprot.org/citations/19158675" target="\_blank">19158675</a>, PubMed:<a href="http://www.uniprot.org/citations/19158676" target="\_blank">19158676</a>, PubMed:<a href="http://www.uniprot.org/citations/19234215" target="\_blank">19234215</a>, PubMed:<a href="http://www.uniprot.org/citations/19494289" target="\_blank">19494289</a>, PubMed:<a href="http://www.uniprot.org/citations/21487011" target="\_blank">21487011</a>, PubMed:<a href="http://www.uniprot.org/citations/24630722" target="\_blank">24630722</a>, PubMed:<a href="http://www.uniprot.org/citations/25847972" target="\_blank">25847972</a>, PubMed:<a href="http://www.uniprot.org/citations/30674671" target="\_blank">30674671</a>, PubMed:<a href="http://www.uniprot.org/citations/34678144" target="\_blank">34678144</a>, PubMed:<a href="http://www.uniprot.org/citations/36050480" target="\_blank">36050480</a>). Promotes caspase- mediated apoptosis involving predominantly caspase-8 and also caspase-9 in a probable cell type-specific manner (PubMed:<a href="http://www.uniprot.org/citations/11103777" target="\_blank">11103777</a>, PubMed:<a href="http://www.uniprot.org/citations/12646168" target="\_blank">12646168</a>). Involved in activation of the mitochondrial apoptotic pathway, promotes caspase-8-dependent proteolytic maturation of BID independently of FADD in certain cell types and also mediates mitochondrial translocation of BAX and activates BAX-dependent apoptosis coupled to activation of caspase-9, -2 and -3 (PubMed:<a href="http://www.uniprot.org/citations/14730312" target="\_blank">14730312</a>, PubMed:<a href="http://www.uniprot.org/citations/16964285" target="\_blank">16964285</a>). Involved in innate immune response by acting as an integral adapter in the assembly of various inflammasomes (NLRP1, NLRP2, NLRP3, NLRP6, AIM2 and probably IFI16) which recruit and activate caspase-1 leading to processing and secretion of pro-inflammatory cytokines (PubMed:<a href="http://www.uniprot.org/citations/15030775" target="\_blank">15030775</a>, PubMed:<a href="http://www.uniprot.org/citations/16982856" target="\_blank">16982856</a>, PubMed:<a href="http://www.uniprot.org/citations/17349957" target="\_blank">17349957</a>, PubMed:<a href="http://www.uniprot.org/citations/17599095" target="\_blank">17599095</a>, PubMed:<a href="http://www.uniprot.org/citations/19158675" target="\_blank">19158675</a>, PubMed:<a href="http://www.uniprot.org/citations/19158676" target="\_blank">19158676</a>, PubMed:<a href="http://www.uniprot.org/citations/19234215" target="\_blank">19234215</a>, PubMed:<a href="http://www.uniprot.org/citations/21487011" target="\_blank">21487011</a>, PubMed:<a href="http://www.uniprot.org/citations/23530044" target="\_blank">23530044</a>, PubMed:<a href="http://www.uniprot.org/citations/24630722" target="\_blank">24630722</a>, PubMed:<a href="http://www.uniprot.org/citations/25847972" target="\_blank">25847972</a>, PubMed:<a href="http://www.uniprot.org/citations/29440442" target="\_blank">29440442</a>, PubMed:<a href="http://www.uniprot.org/citations/30674671" target="\_blank">30674671</a>, PubMed:<a href="http://www.uniprot.org/citations/33980849" target="\_blank">33980849</a>, PubMed:<a href="http://www.uniprot.org/citations/34678144" target="\_blank">34678144</a>, PubMed:<a href="http://www.uniprot.org/citations/34706239" target="\_blank">34706239</a>). Caspase-1-dependent inflammation leads to macrophage pyroptosis, a form of cell death (PubMed:<a href="http://www.uniprot.org/citations/24630722" target="\_blank">24630722</a>). The function as activating adapter in different types of inflammasomes is mediated by the pyrin and CARD domains and their homotypic interactions (PubMed:<a href="http://www.uniprot.org/citations/14499617" target="\_blank">14499617</a>, PubMed:<a href="http://www.uniprot.org/citations/19234215" target="\_blank">19234215</a>, PubMed:<a href="http://www.uniprot.org/citations/24630722" target="\_blank">24630722</a>). Clustered PYCARD nucleates the formation of caspase-1 filaments through the interaction of their respective CARD domains, acting as a platform for of caspase-1 polymerization (PubMed:<a href="http://www.uniprot.org/citations/14499617" target="\_blank">14499617</a>, PubMed:<a href="http://www.uniprot.org/citations/19234215" target="\_blank">19234215</a>, PubMed:<a href="http://www.uniprot.org/citations/24630722" target="\_blank">24630722</a>).

<http://www.uniprot.org/citations/24630722> target="\_blank">24630722</a>). In the NLRP1 and NLRC4 inflammasomes seems not be required but facilitates the processing of procaspase-1 (PubMed:<a href="http://www.uniprot.org/citations/17349957" target="\_blank">17349957</a>). In cooperation with NOD2 involved in an inflammasome activated by bacterial muramyl dipeptide leading to caspase-1 activation (PubMed:<a href="http://www.uniprot.org/citations/16964285" target="\_blank">16964285</a>). May be involved in RIGI-triggered pro-inflammatory responses and inflammasome activation (PubMed:<a href="http://www.uniprot.org/citations/19915568" target="\_blank">19915568</a>). In collaboration with AIM2 which detects cytosolic double-stranded DNA may also be involved in a caspase-1-independent cell death that involves caspase-8 (PubMed:<a href="http://www.uniprot.org/citations/19158675" target="\_blank">19158675</a>, PubMed:<a href="http://www.uniprot.org/citations/19158676" target="\_blank">19158676</a>). In adaptive immunity may be involved in maturation of dendritic cells to stimulate T-cell immunity and in cytoskeletal rearrangements coupled to chemotaxis and antigen uptake may be involved in post-transcriptional regulation of the guanine nucleotide exchange factor DOCK2; the latter function is proposed to involve the nuclear form (PubMed:<a href="http://www.uniprot.org/citations/22732093" target="\_blank">22732093</a>). Also involved in transcriptional activation of cytokines and chemokines independent of the inflammasome; this function may involve AP-1, NF-kappa-B, MAPK and caspase-8 signaling pathways (PubMed:<a href="http://www.uniprot.org/citations/12486103" target="\_blank">12486103</a>, PubMed:<a href="http://www.uniprot.org/citations/16585594" target="\_blank">16585594</a>). For regulation of NF-kappa-B activating and inhibiting functions have been reported (PubMed:<a href="http://www.uniprot.org/citations/12486103" target="\_blank">12486103</a>). Modulates NF-kappa-B induction at the level of the IKK complex by inhibiting kinase activity of CHUK and IKBK (PubMed:<a href="http://www.uniprot.org/citations/12486103" target="\_blank">12486103</a>, PubMed:<a href="http://www.uniprot.org/citations/16585594" target="\_blank">16585594</a>). Proposed to compete with RIPK2 for association with CASP1 thereby down-regulating CASP1-mediated RIPK2-dependent NF-kappa-B activation and activating interleukin-1 beta processing (PubMed:<a href="http://www.uniprot.org/citations/16585594" target="\_blank">16585594</a>). Modulates host resistance to DNA virus infection, probably by inducing the cleavage of and inactivating CGAS in presence of cytoplasmic double-stranded DNA (PubMed:<a href="http://www.uniprot.org/citations/28314590" target="\_blank">28314590</a>).

### Cellular Location

Cytoplasm. Inflammasome. Endoplasmic reticulum. Mitochondrion. Nucleus Note=Upstream of caspase activation, a redistribution from the cytoplasm to the aggregates occurs. These appear as hollow, perinuclear spherical, ball-like structures (PubMed:11103777, PubMed:12191486, PubMed:15030775). Upon NLRP3 inflammasome activation redistributes to the perinuclear space localizing to endoplasmic reticulum and mitochondria (PubMed:12191486, PubMed:15030775). Localized primarily to the nucleus in resting monocytes/macrophages and rapidly redistributed to the cytoplasm upon pathogen infection (PubMed:19234215). Localized to large cytoplasmic aggregate appearing as a speck containing AIM2, PYCARD, CASP8 and bacterial DNA after infection with Francisella tularensis (By similarity). {ECO:0000250|UniProtKB:Q9EPB4, ECO:0000269|PubMed:11103777, ECO:0000269|PubMed:12191486, ECO:0000269|PubMed:15030775, ECO:0000269|PubMed:19234215}

### Tissue Location


Widely expressed at low levels. Detected in peripheral blood leukocytes, lung, small intestine, spleen, thymus, colon and at lower levels in placenta, liver and kidney. Very low expression in skeletal muscle, heart and brain. Expressed in lung epithelial cells (at protein level) (PubMed:23229815). Detected in the leukemia cell lines HL-60 and U-937, but not in Jurkat T-cell lymphoma and Daudi Burkitt's lymphoma. Detected in the melanoma cell line WM35, but not in WM793. Not detected in HeLa cervical carcinoma cells and MOLT-4 lymphocytic leukemia cells.

### Mouse Monoclonal Antibody to PYCARD - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**Mouse Monoclonal Antibody to PYCARD - Images**




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Western blot analysis using PYCARD mAb against human PYCARD (AA: 1-120) recombinant protein. (Expected MW is 39.2 kDa)



Western blot analysis using PYCARD mAb against HEK2993 (1) and PYCARD (AA: 1-120)-hlgGfc transfected HEK2993 (2) cell lysate.




Flow cytometric analysis of HEK2993 cells using PYCARD mouse mAb (green) and negative control (red).




Immunofluorescence analysis of GC-7901 cells using PYCARD mouse mAb (green). Blue: DRAQ5



fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor- 555 phalloidin. Secondary antibody from Fisher



Immunofluorescence analysis of HeLa cells using PYCARD mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor- 555 phalloidin. Secondary antibody from Fisher



Immunohistochemical analysis of paraffin-embedded ovarian cancer tissues using PYCARD mouse mAb with DAB staining.

#### **Mouse Monoclonal Antibody to PYCARD - References**

1.J Invest Dermatol. 2013 Feb;133(2):518-27. ; 2.Cancer Sci. 2010 Aug;101(8):1822-7.;

#### **Mouse Monoclonal Antibody to PYCARD - Citations**

- [Promoter methylation status of tumor suppressor genes and inhibition of expression of DNA methyltransferase 1 in non-small cell lung cancer.](#)