

**Mouse Monoclonal Antibody to C3C**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO2388a**

**Specification**

---

**Mouse Monoclonal Antibody to C3C - Product Information**

Application	E, WB, IHC
Primary Accession	<a href="#">P01024</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1
Calculated MW	187.1kDa KDa

**Description**

Complement component C3 plays a central role in the activation of complement system. Its activation is required for both classical and alternative complement activation pathways. The encoded preproprotein is proteolytically processed to generate alpha and beta subunits that form the mature protein, which is then further processed to generate numerous peptide products. The C3a peptide, also known as the C3a anaphylatoxin, modulates inflammation and possesses antimicrobial activity. Mutations in this gene are associated with atypical hemolytic uremic syndrome and age-related macular degeneration in human patients.;

**Immunogen**

Purified recombinant fragment of human C3C (AA: 1521-1649) expressed in E. Coli.

**Formulation**

Purified antibody in PBS with 0.05% sodium azide

**Application Note**

ELISA: 1/10000; WB: 1/500 - 1/2000; IHC: 1/200 - 1/1000;

**Mouse Monoclonal Antibody to C3C - Additional Information**

**Gene ID** 718

**Other Names**

ASP; C3a; C3b; AHUS5; ARMD9; CPAMD1; HEL-S-62p

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Mouse Monoclonal Antibody to C3C is for research use only and not for use in diagnostic or therapeutic procedures.

**Mouse Monoclonal Antibody to C3C - Protein Information**

**Name** C3

**Synonyms** CPAMD1

**Function**

C3 plays a central role in the activation of the complement system. Its processing by C3 convertase is the central reaction in both classical and alternative complement pathways. After activation C3b can bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates. [C3-beta-c]: Acts as a chemoattractant for neutrophils in chronic inflammation.

**Cellular Location**

Secreted.

**Tissue Location**


Plasma. The acylation stimulating protein (ASP) is expressed in adipocytes and released into the plasma during both the fasting and postprandial periods.

**Mouse Monoclonal Antibody to C3C - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**Mouse Monoclonal Antibody to C3C - Images**




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Western blot analysis using C3C mAb against human C3C (AA: 1521-1649) recombinant protein. (Expected MW is 40.7 kDa)



Western blot analysis using C3C mAb against HEK293 (1) and C3C (AA: 1521-1649)-hIgGfC transfected HEK293 (2) cell lysate.



Immunohistochemical analysis of paraffin-embedded ovarian cancer tissues using C3C mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded endometrial cancer tissues using C3C mouse mAb with DAB staining.

#### **Mouse Monoclonal Antibody to C3C - References**

1.Blood. 2015 Apr 9;125(15):2359-69. ; 2.Mediators Inflamm. 2013;2013:716902.;