

Mouse Monoclonal Antibody to ESRRA
Purified Mouse Monoclonal Antibody
Catalog # AO2421a**Specification**

Mouse Monoclonal Antibody to ESRRA - Product Information

Application	E, WB, FC
Primary Accession	P11474
Reactivity	Human, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1
Calculated MW	45.5kDa KDa

Description

The protein encoded by this gene is a nuclear receptor that is closely related to the estrogen receptor. This protein acts as a site-specific transcription regulator and has been also shown to interact with estrogen and the transcription factor TFIIB by direct protein-protein contact. The binding and regulatory activities of this protein have been demonstrated in the regulation of a variety of genes including lactoferrin, osteopontin, medium-chain acyl coenzyme A dehydrogenase (MCAD) and thyroid hormone receptor genes. A processed pseudogene of ESRRA is located on chromosome 13q12.1. Alternatively spliced transcript variants have been found for this gene.;

Immunogen

Purified recombinant fragment of human ESRRA (AA: 198-376) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

Application Note

ELISA: 1/10000; WB: 1/500 - 1/2000; FCM: 1/200 - 1/400

Mouse Monoclonal Antibody to ESRRA - Additional Information

Gene ID 2101

Other Names

ERR1; ERRa; ESRL1; NR3B1; ERRalpha

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Monoclonal Antibody to ESRRA is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Monoclonal Antibody to ESRRA - Protein Information

Name ESRRRA

Synonyms ERR1, ESRL1, NR3B1

Function

Binds to an ERR-alpha response element (ERRE) containing a single consensus half-site, 5'-TNAAGGTCA-3'. Can bind to the medium- chain acyl coenzyme A dehydrogenase (MCAD) response element NRRE-1 and may act as an important regulator of MCAD promoter. Binds to the C1 region of the lactoferrin gene promoter. Requires dimerization and the coactivator, PGC-1A, for full activity. The ERRalpha/PGC1alpha complex is a regulator of energy metabolism. Induces the expression of PERM1 in the skeletal muscle.

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00407, ECO:0000269|PubMed:18063693, ECO:0000269|PubMed:21190936}. Cytoplasm. Note=Co-localizes to the cytoplasm only in presence of MAPK15.

Mouse Monoclonal Antibody to ESRRRA - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Mouse Monoclonal Antibody to ESRRRA - Images




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Western blot analysis using ESRR mAb against human ESRR (AA: 198-376) recombinant protein. (Expected MW is 45.3 kDa)



Western blot analysis using ESRR mAb against HEK293 (1) and ESRR (AA: 198-376)-hIgGfC transfected HEK293 (2) cell lysate.



Western blot analysis using ESRRRA mouse mAb against C6 (1) cell lysate.



Flow cytometric analysis of Hela cells using ESRRRA mouse mAb (green) and negative control (red).

Mouse Monoclonal Antibody to ESRRRA - References

1. J Huazhong Univ Sci Technolog Med Sci. 2014 Dec;34(6):875-81. ; 2. Mol Ther. 2014 Apr;22(4):743-51. ;