

#### **HAVCR1**

Purified Mouse Monoclonal Antibody Catalog # AO2576a

# **Specification**

### **HAVCR1 - Product Information**

Application E, WB
Primary Accession Q96D42
Reactivity Human
Host Mouse
Clonality Monoclonal
Isotype Mouse IgG1
Calculated MW 38.7kDa KDa

**Immunogen** 

Purified recombinant fragment of human HAVCR1 (AA: 70-290) expressed in E. Coli.

#### **Formulation**

Purified antibody in PBS with 0.05% sodium azide

#### **HAVCR1 - Additional Information**

**Gene ID 26762** 

#### **Other Names**

TIM; KIM1; TIM1; CD365; HAVCR; KIM-1; TIM-1; TIMD1; TIMD-1; HAVCR-1

### **Dilution**

E~~ 1/10000

WB~~ 1/500 - 1/2000

# **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

HAVCR1 is for research use only and not for use in diagnostic or therapeutic procedures.

### **HAVCR1 - Protein Information**

#### Name HAVCR1

Synonyms KIM1, TIM1, TIMD1

# **Function**

Phosphatidylserine receptor that plays an important functional role in regulatory B-cells homeostasis including generation, expansion and suppressor functions (By similarity). As P-selectin/SELPLG ligand, plays a specialized role in activated but not naive T-cell trafficking during



inflammatory responses (PubMed:<a href="http://www.uniprot.org/citations/24703780" target="\_blank">24703780</a>). Controls thereby T-cell accumulation in the inflamed central nervous system (CNS) and the induction of autoimmune disease (PubMed:<a href="http://www.uniprot.org/citations/24703780" target="\_blank">24703780</a>). Regulates also expression of various anti- inflammatory cytokines and co-inhibitory ligands including IL10 (By similarity). Acts as a regulator of T-cell proliferation (By similarity). May play a role in kidney injury and repair (PubMed:<a href="http://www.uniprot.org/citations/17471468" target=" blank">17471468</a>).

# **Cellular Location**

Cell membrane; Single-pass type I membrane protein

#### **Tissue Location**

Widely expressed, with highest levels in kidney and testis. Expressed by activated CD4+ T-cells during the development of helper T-cells responses.

### **HAVCR1 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# **HAVCR1** - Images




Figure 1:Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)






Figure 2:Western blot analysis using HAVCR1 mAb against human HAVCR1 (AA: 70-290) recombinant protein. (Expected MW is 37 kDa)




Figure 3:Western blot analysis using HAVCR1 mAb against HEK293 (1) and HAVCR1 (AA: 70-290)-hlgGFc transfected HEK293 (2) cell lysate.




Figure 4:Flow cytometric analysis of Hela cells using HAVCR1 mouse mAb (green) and negative





control (red).

# **HAVCR1 - References**

1.Biomed Res Int. 2015;2015:854070. 2.Pediatr Res. 2015 Oct;78(4):430-5.