

BCL6
Purified Mouse Monoclonal Antibody
Catalog # AO2648a

Specification

BCL6 - Product Information

Application	E, WB
Primary Accession	P41182
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1
Calculated MW	78.8kDa KDa

Immunogen

Purified recombinant fragment of human BCL6 (AA: 147-276) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

BCL6 - Additional Information

Gene ID 604

Other Names

BCL5; LAZ3; BCL6A; ZNF51; ZBTB27

Dilution

E~~ 1/10000
WB~~ 1/500 - 1/2000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BCL6 is for research use only and not for use in diagnostic or therapeutic procedures.

BCL6 - Protein Information

Name BCL6

Synonyms BCL5, LAZ3, ZBTB27, ZNF51

Function

Transcriptional repressor mainly required for germinal center (GC) formation and antibody affinity maturation which has different mechanisms of action specific to the lineage and biological functions. Forms complexes with different corepressors and histone deacetylases to repress the

transcriptional expression of different subsets of target genes. Represses its target genes by binding directly to the DNA sequence 5'-TTCCTAGAA-3' (BCL6-binding site) or indirectly by repressing the transcriptional activity of transcription factors. In GC B-cells, represses genes that function in differentiation, inflammation, apoptosis and cell cycle control, also autoregulates its transcriptional expression and up-regulates, indirectly, the expression of some genes important for GC reactions, such as AICDA, through the repression of microRNAs expression, like miR155. An important function is to allow GC B-cells to proliferate very rapidly in response to T- cell dependent antigens and tolerate the physiological DNA breaks required for immunoglobulin class switch recombination and somatic hypermutation without inducing a p53/TP53-dependent apoptotic response. In follicular helper CD4(+) T-cells (T(FH) cells), promotes the expression of T(FH)-related genes but inhibits the differentiation of T(H)1, T(H)2 and T(H)17 cells. Also required for the establishment and maintenance of immunological memory for both T- and B-cells. Suppresses macrophage proliferation through competition with STAT5 for STAT- binding motifs binding on certain target genes, such as CCL2 and CCND2. In response to genotoxic stress, controls cell cycle arrest in GC B- cells in both p53/TP53-dependendent and -independent manners. Besides, also controls neurogenesis through the alteration of the composition of NOTCH-dependent transcriptional complexes at selective NOTCH targets, such as HES5, including the recruitment of the deacetylase SIRT1 and resulting in an epigenetic silencing leading to neuronal differentiation.

Cellular Location

Nucleus

Tissue Location

Expressed in germinal center T- and B-cells and in primary immature dendritic cells.

BCL6 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BCL6 - Images




Figure 1: Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)




Figure 3: Western blot analysis using BCL6 mAb against HEK293 (1) and BCL6 (AA: 147-276)-hlgGfc transfected HEK293 (2) cell lysate.




Figure 3: Western blot analysis using BCL6 mAb against human BCL6 (AA: 147-276) recombinant protein. (Expected MW is 40.5 kDa)

BCL6 - References

1. Cancer Lett. 2015 Sep 1;365(2):190-200.
2. BMC Cancer. 2014 Jun 10;14:418.