

**CADM1 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP10703a****Specification**

---

**CADM1 Antibody (N-term) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">Q9BY67</a>
Other Accession	<a href="#">Q8R5M8</a> , <a href="#">NP_001091987.1</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	48509
Antigen Region	66-94

**CADM1 Antibody (N-term) - Additional Information****Gene ID** 23705**Other Names**

Cell adhesion molecule 1, Immunoglobulin superfamily member 4, IgSF4, Nectin-like protein 2, NECL-2, Spermatogenic immunoglobulin superfamily, SgIgSF, Synaptic cell adhesion molecule, SynCAM, Tumor suppressor in lung cancer 1, TSLC-1, CADM1 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=5951](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=5951))

**Target/Specificity**

This CADM1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 66-94 amino acids from the N-terminal region of human CADM1.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CADM1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CADM1 Antibody (N-term) - Protein Information**

**Name** CADM1 ([HGNC:5951](#))

**Function** Mediates homophilic cell-cell adhesion in a  $\text{Ca}^{2+}$ - independent manner (PubMed:[12050160](#), PubMed:[22438059](#)). Also mediates heterophilic cell-cell adhesion with CADM3 and NECTIN3 in a  $\text{Ca}^{2+}$ - independent manner (By similarity). Interaction with CRTAM promotes natural killer (NK) cell cytotoxicity and interferon-gamma (IFN-gamma) secretion by CD8+ cells in vitro as well as NK cell-mediated rejection of tumors expressing CADM1 in vivo (PubMed:[15811952](#)). In mast cells, may mediate attachment to and promote communication with nerves (PubMed:[15905536](#)). CADM1, together with MITF, is essential for development and survival of mast cells in vivo (PubMed:[22438059](#)). By interacting with CRTAM and thus promoting the adhesion between CD8+ T- cells and CD8+ dendritic cells, regulates the retention of activated CD8+ T-cell within the draining lymph node (By similarity). Required for the intestinal retention of intraepithelial CD4+ CD8+ T-cells and, to a lesser extent, intraepithelial and lamina propria CD8+ T-cells and CD4+ T-cells (By similarity). Interaction with CRTAM promotes the adhesion to gut-associated CD103+ dendritic cells, which may facilitate the expression of gut-homing and adhesion molecules on T-cells and the conversion of CD4+ T-cells into CD4+ CD8+ T-cells (By similarity). Acts as a synaptic cell adhesion molecule and plays a role in the formation of dendritic spines and in synapse assembly (By similarity). May be involved in neuronal migration, axon growth, pathfinding, and fasciculation on the axons of differentiating neurons (By similarity). May play diverse roles in the spermatogenesis including in the adhesion of spermatocytes and spermatids to Sertoli cells and for their normal differentiation into mature spermatozoa (By similarity). Acts as a tumor suppressor in non-small-cell lung cancer (NSCLC) cells (PubMed:[11279526](#), PubMed:[12234973](#)). May contribute to the less invasive phenotypes of lepidic growth tumor cells (PubMed:[12920246](#)).


**Cellular Location**


Cell membrane; Single-pass type I membrane protein. Synapse  
{ECO:0000250|UniProtKB:Q8R5M8} Note=Localized to the basolateral plasma membrane of epithelial cells in gall bladder. {ECO:0000250|UniProtKB:Q8R5M8}

**CADM1 Antibody (N-term) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**CADM1 Antibody (N-term) - Images**



CADM1 Antibody (N-term) (Cat. #AP10703a) western blot analysis in mouse cerebellum tissue lysates (35ug/lane). This demonstrates the CADM1 antibody detected the CADM1 protein (arrow).



CADM1 Antibody (N-term) (Cat. #AP10703a) western blot analysis in MDA-MB435 cell line lysates (35ug/lane). This demonstrates the CADM1 antibody detected the CADM1 protein (arrow).



CADM1 antibody (N-term) (Cat. #AP10703a) immunohistochemistry analysis in formalin fixed and paraffin embedded human lung carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CADM1 antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

#### **CADM1 Antibody (N-term) - Background**

Mediates homophilic cell-cell adhesion in a  $\text{Ca}^{2+}$ -independent manner. Also mediates heterophilic cell-cell adhesion with CADM3 and PVRL3 in a  $\text{Ca}^{2+}$ -independent manner. Acts as a

tumor suppressor in non-small-cell lung cancer (NSCLC) cells. Interaction with CRTAM promotes natural killer (NK) cell cytotoxicity and interferon-gamma (IFN-gamma) secretion by CD8+ cells in vitro as well as NK cell-mediated rejection of tumors expressing CADM3 in vivo. May contribute to the less invasive phenotypes of lepidic growth tumor cells. In mast cells, may mediate attachment to and promote communication with nerves. CADM1, together with MITF, is essential for development and survival of mast cells in vivo. May act as a synaptic cell adhesion molecule that drives synapse assembly. May be involved in neuronal migration, axon growth, pathfinding, and fasciculation on the axons of differentiating neurons. May play diverse roles in the spermatogenesis including in the adhesion of spermatocytes and spermatids to Sertoli cells and for their normal differentiation into mature spermatozoa.

#### **CADM1 Antibody (N-term) - References**

Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)  
Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)  
McGovern, D.P., et al. Hum. Mol. Genet. 19(17):3468-3476(2010)  
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :  
You, Y., et al. Melanoma Res. 20(3):179-183(2010)