

**RNT2 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP10764a**

**Specification**

---

**RNT2 Antibody (N-term) - Product Information**

Application	<b>WB, IHC-P,E</b>
Primary Accession	<a href="#">O00584</a>
Other Accession	<a href="#">NP_003721.2</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Antigen Region	<b>25-55</b>

**RNT2 Antibody (N-term) - Additional Information**

**Gene ID** 8635

**Other Names**

Ribonuclease T2, 3127-, Ribonuclease 6, RNASET2, RNASE6PL

**Target/Specificity**

This RNT2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 25-55 amino acids from the N-terminal region of human RNT2.

**Dilution**

WB~~1:500-1:1000

IHC-P~~1:50~100

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

RNT2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**RNT2 Antibody (N-term) - Protein Information**

**Name** RNASET2

**Synonyms** RNASE6PL

**Function** Ribonuclease that plays an essential role in innate immune response by recognizing and degrading RNAs from microbial pathogens that are subsequently sensed by TLR8 (PubMed:[31778653](#)). Cleaves preferentially single-stranded RNA molecules between purine and uridine residues, which critically contributes to the supply of catabolic uridine and the generation of purine-2',3'-cyclophosphate-terminated oligoribonucleotides (PubMed:[31778653](#)). In turn, RNase T2 degradation products promote the RNA-dependent activation of TLR8 (PubMed:[31778653](#)). Also plays a key role in degradation of mitochondrial RNA and processing of non-coding RNA imported from the cytosol into mitochondria (PubMed:[28730546](#), PubMed:[30184494](#)). Participates as well in degradation of mitochondrion-associated cytosolic rRNAs (PubMed:[30385512](#)).

#### Cellular Location

Secreted. Lysosome lumen. Endoplasmic reticulum lumen. Mitochondrion intermembrane space. Note=Full-length RNASET2 is found in the endoplasmic reticulum while smaller RNASET2 proteolytic products are found in the lysosome fraction.

#### Tissue Location


Ubiquitous. Higher expression levels observed in the temporal lobe and fetal brain.

### RNT2 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### RNT2 Antibody (N-term) - Images



Western blot analysis of lysates from 293T cell line, human brain tissue lysate (from left to right), using RNT2 Antibody (N-term)(Cat. #AP10764a). AP10764a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.



RNT2 antibody (N-term) (Cat. #AP10764a) immunohistochemistry analysis in formalin fixed and paraffin embedded human spleen carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the RNT2 antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

#### **RNT2 Antibody (N-term) - Background**

This ribonuclease gene is a novel member of the Rh/T2/S-glycoprotein class of extracellular ribonucleases. It is a single copy gene that maps to 6q27, a region associated with human malignancies and chromosomal rearrangement.

#### **RNT2 Antibody (N-term) - References**

Fransen, K., et al. Hum. Mol. Genet. 19(17):3482-3488(2010)  
Quan, C., et al. Nat. Genet. 42(7):614-618(2010)  
Liu, J., et al. J. Cell. Mol. Med. 14(5):1146-1155(2010)  
Henneke, M., et al. Nat. Genet. 41(7):773-775(2009)  
Monti, L., et al. Oncol. Res. 17(2):69-74(2008)

#### **RNT2 Antibody (N-term) - Citations**

- [Regulation of mitochondrion-associated cytosolic ribosomes by mammalian mitochondrial ribonuclease T2 \(RNASET2\).](#)