

ENT1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1086C

Specification

ENT1 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O99808
Other Accession	O54698
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	234-263

ENT1 Antibody (Center) - Additional Information

Gene ID 2030

Other Names

Equilibrative nucleoside transporter 1, Equilibrative nitrobenzylmercaptapurine riboside-sensitive nucleoside transporter, Equilibrative NBMPR-sensitive nucleoside transporter, Nucleoside transporter, es-type, Solute carrier family 29 member 1, SLC29A1, ENT1

Target/Specificity

This ENT1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 234-263 amino acids from the Central region of rat ENT1(O54698).

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ENT1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ENT1 Antibody (Center) - Protein Information

Name SLC29A1 ([HGNC:11003](#))

Synonyms ENT1

Function Uniporter involved in the facilitative transport of nucleosides and nucleobases, and contributes to maintaining their cellular homeostasis (PubMed:[10722669](#), PubMed:[10755314](#), PubMed:[12527552](#), PubMed:[14759222](#), PubMed:[15037197](#), PubMed:[17379602](#), PubMed:[21795683](#), PubMed:[26406980](#), PubMed:[27995448](#), PubMed:[35790189](#), PubMed:[8986748](#)). Functions as a Na(+)-independent transporter (PubMed:[8986748](#)). Involved in the transport of nucleosides such as adenosine, guanosine, inosine, uridine, thymidine and cytidine (PubMed:[10722669](#), PubMed:[10755314](#), PubMed:[12527552](#), PubMed:[14759222](#), PubMed:[15037197](#), PubMed:[17379602](#), PubMed:[26406980](#), PubMed:[8986748](#)). Also transports purine nucleobases (hypoxanthine, adenine, guanine) and pyrimidine nucleobases (thymine, uracil) (PubMed:[21795683](#), PubMed:[27995448](#)). Mediates basolateral nucleoside uptake into Sertoli cells, thereby regulating the transport of nucleosides in testis across the blood-testis barrier (By similarity). Regulates inosine levels in brown adipocytes tissues (BAT) and extracellular inosine levels, which controls BAT-dependent energy expenditure (PubMed:[35790189](#)).

Cellular Location

Basolateral cell membrane; Multi-pass membrane protein. Apical cell membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Note=Localized to the basolateral membrane of Sertoli cells (PubMed:23639800). Localized to the cell membrane of erythrocytes (PubMed:11584005, PubMed:23219802).

Tissue Location


Expressed in testis at the blood-testis barrier (at protein level) (PubMed:23639800). Detected in erythrocytes (at protein level) (PubMed:11584005, PubMed:23219802). Expressed at relatively high levels in cerebral cortex, particularly the frontal and parietal lobes, and the thalamus and basal ganglia (at protein level) (PubMed:11311901). In the midbrain expressed at moderate levels, whereas in the other areas of the brainstem, namely medulla and pons, cerebellum and the hippocampus expressed at lower amounts when compared to the other brain regions (at protein level) (PubMed:11311901) Expressed in Langerhans cells and lymphocytes in the pancreas (at protein level) (PubMed:15501974). Expressed in kidney, in polarized renal epithelial cells (PubMed:12527552). Expressed in adipose tissues (PubMed:35790189). Expressed in placenta (PubMed:8986748). Expressed in small intestine (PubMed:10755314).


ENT1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ENT1 Antibody (Center) - Images





Western blot analysis of lysate from HL-60 cell line, using ENT1(Slc29a1) Antibody (Center)(Cat. #AP1086c). AP1086c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.



Anti-ENT1(Slc29a1) Antibody (Center) at 1:2000 dilution + mouse lung lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 50 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

ENT1 Antibody (Center) - Background

ENT1 is a member of the equilibrative nucleoside transporter family. It is a transmembrane glycoprotein that localizes to the plasma and mitochondrial membranes and mediates the cellular uptake of nucleosides from the surrounding medium. The protein is categorized as an equilibrative (as opposed to concentrative) transporter that is sensitive to inhibition by nitrobenzylthioinosine (NBMPR). Nucleoside transporters are required for nucleotide synthesis in cells that lack de novo nucleoside synthesis pathways, and are also necessary for the uptake of cytotoxic nucleosides used for cancer and viral chemotherapies.

ENT1 Antibody (Center) - References

- Bone,D.B.,Am. J. Physiol. Heart Circ. Physiol. 293 (6), H3325-H3332 (2007)
- Damaraju,V.L., Am. J. Physiol. Renal Physiol. 293 (1), F200-F211 (2007)

Abdulla,P.,Nucleosides Nucleotides Nucleic Acids 26 (1), 99-110 (2007)

Sundaram,M., J. Biol. Chem. 276 (48), 45270-45275 (2001)

ENT1 Antibody (Center) - Citations

- [Reduced ribavirin antiviral efficacy via nucleoside transporter-mediated drug resistance.](#)