

OR6C4 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10908b

Specification

OR6C4 Antibody (C-term) - Product Information

Application	WB, FC,E
Primary Accession	O8NGE1
Other Accession	NP_001005494.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35010
Antigen Region	242-270

OR6C4 Antibody (C-term) - Additional Information

Gene ID 341418

Other Names

Olfactory receptor 6C4, Olfactory receptor OR12-10, OR6C4

Target/Specificity

This OR6C4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 242-270 amino acids from the C-terminal region of human OR6C4.

Dilution

WB~~1:1000

FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR6C4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR6C4 Antibody (C-term) - Protein Information

Name OR6C4

Function Odorant receptor.


Cellular Location

Cell membrane; Multi-pass membrane protein.


OR6C4 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR6C4 Antibody (C-term) - Images

All lanes : Anti-OR6C4 Antibody (C-term) at 1:1000 dilution Lane 1: MCF-7 whole cell lysate Lane 2: Y79 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 35 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



OR6C4 Antibody (C-term) (Cat. #AP10908b) flow cytometric analysis of MCF-7 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

OR6C4 Antibody (C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR6C4 Antibody (C-term) - References

- Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)
- Youngblood, K., et al. J. Clin. Invest. 93(2):852-861(1994)
- Lauzurica, P., et al. J. Immunol. 148(11):3624-3630(1992)