

RAP1GDS1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11945c

Specification

RAP1GDS1 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	P52306
Other Accession	NP_001093900.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	66317
Antigen Region	217-244

RAP1GDS1 Antibody (Center) - Additional Information

Gene ID 5910

Other Names

Rap1 GTPase-GDP dissociation stimulator 1, Exchange factor smgGDS, SMG GDS protein, SMG P21 stimulatory GDP/GTP exchange protein, RAP1GDS1

Target/Specificity

This RAP1GDS1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 217-244 amino acids from the Central region of human RAP1GDS1.

Dilution

WB~~1:1000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RAP1GDS1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

RAP1GDS1 Antibody (Center) - Protein Information

Name RAP1GDS1 {ECO:0000303|Ref.3}

Synonyms SMGGDS {ECO:0000303|Ref.3}

Function Acts as a GEF (guanine nucleotide exchange factor) for the Rho family of small GTP-binding proteins (G proteins) that stimulates the dissociation of GDP to enable subsequent binding of GTP (PubMed:[11948427](#), PubMed:[1549351](#), PubMed:[20709748](#), PubMed:[28630045](#), PubMed:[30190425](#)). Additionally, appears to chaperone the processing and/or trafficking of small GTPases containing a C-terminal polybasic region independently of GEF activity (PubMed:[20709748](#), PubMed:[21242305](#)). Targets include RAP1A/RAP1B, RHOA, RHOB, RHOC, RAC1 and KRAS (PubMed:[11948427](#), PubMed:[1549351](#), PubMed:[20709748](#), PubMed:[24415755](#)). Regulates mitochondrial dynamics by controlling RHOT function to promote mitochondrial fission during high calcium conditions (PubMed:[27716788](#)). Able to promote the Ca(2+) release from the endoplasmic reticulum via both inositol trisphosphate (Ins3P) and ryanodine sensitive receptors leading to a enhanced mitochondrial Ca(2+) uptake (PubMed:[24349085](#)).

Cellular Location


Cytoplasm, cytosol. Endoplasmic reticulum. Mitochondrion. Nucleus Note=Nuclear import is dependent on complexing with a GTPase containing a C-terminal polybasic region.

RAP1GDS1 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RAP1GDS1 Antibody (Center) - Images



RAP1GDS1 Antibody (Center) (Cat. #AP11945c) western blot analysis in HepG2,MDA-MB231 cell line lysates (35ug/lane).This demonstrates the RAP1GDS1 antibody detected the RAP1GDS1 protein (arrow).



RAP1GDS1 Antibody (Center) (Cat. #AP11945c) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of RAP1GDS1 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

RAP1GDS1 Antibody (Center) - Background

The smg GDP dissociation stimulator (smgGDS) protein is a stimulatory GDP/GTP exchange protein with GTPase activity (Riess et al., 1993 [PubMed 8262526]).

RAP1GDS1 Antibody (Center) - References

- Schönherr, C., et al. *Oncogene* 29(19):2817-2830(2010)
- Branham, M.T., et al. *J. Biol. Chem.* 284(37):24825-24839(2009)
- Zemunik, T., et al. *Croat. Med. J.* 50(1):23-33(2009)
- Tew, G.W., et al. *J. Biol. Chem.* 283(2):963-976(2008)
- Tiwari, S., et al. *Blood* 103(7):2661-2667(2004)