

ATP6V1B1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12290b

Specification

ATP6V1B1 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P15313
Other Accession	NP_001683.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	56833
Antigen Region	472-501

ATP6V1B1 Antibody (C-term) - Additional Information

Gene ID 525

Other Names

V-type proton ATPase subunit B, kidney isoform, V-ATPase subunit B 1, Endomembrane proton pump 58 kDa subunit, Vacuolar proton pump subunit B 1, ATP6V1B1, ATP6B1, VATB, VPP3

Target/Specificity

This ATP6V1B1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 472-501 amino acids from the C-terminal region of human ATP6V1B1.

Dilution

WB~~1:1000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ATP6V1B1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ATP6V1B1 Antibody (C-term) - Protein Information

Name ATP6V1B1

Synonyms ATP6B1, VATB, VPP3

Function Non-catalytic subunit of the V1 complex of vacuolar(H⁺)- ATPase (V-ATPase), a multisubunit enzyme composed of a peripheral complex (V1) that hydrolyzes ATP and a membrane integral complex (V0) that translocates protons (PubMed:[16769747](#)). V-ATPase is responsible for acidifying and maintaining the pH of intracellular compartments and in some cell types, is targeted to the plasma membrane, where it is responsible for acidifying the extracellular environment (PubMed:[32001091](#)). Essential for the proper assembly and activity of V-ATPase (PubMed:[16769747](#)). In renal intercalated cells, mediates secretion of protons (H⁺) into the urine thereby ensuring correct urinary acidification (PubMed:[16769747](#)). Required for optimal olfactory function by mediating the acidification of the nasal olfactory epithelium (By similarity).

Cellular Location

Apical cell membrane. Basolateral cell membrane {ECO:0000250|UniProtKB:Q91YH6}


Tissue Location

Kidney; localizes to early distal nephron, encompassing thick ascending limbs and distal convoluted tubules (at protein level) (PubMed:[16769747](#), PubMed:[29993276](#)). Expressed in the cochlea and endolymphatic sac (PubMed:[9916796](#))


ATP6V1B1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ATP6V1B1 Antibody (C-term) - Images

ATP6V1B1 Antibody (C-term) (Cat. #AP12290b) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the ATP6V1B1 antibody detected the ATP6V1B1 protein (arrow).



ATP6V1B1 Antibody (C-term) (Cat. #AP12290b) immunohistochemistry analysis in formalin fixed and paraffin embedded human breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of ATP6V1B1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

ATP6V1B1 Antibody (C-term) - Background

This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is one of two V1 domain B subunit isoforms and is found in the kidney. Mutations in this gene cause distal renal tubular acidosis associated with sensorineural deafness. [provided by RefSeq].

ATP6V1B1 Antibody (C-term) - References

Bailey, S.D., et al. *Diabetes Care* 33(10):2250-2253(2010)
Sharifian, M., et al. *Iran J Kidney Dis* 4(3):202-206(2010)
Talmud, P.J., et al. *Am. J. Hum. Genet.* 85(5):628-642(2009)
Andreucci, E., et al. *Pediatr. Nephrol.* 24(11):2147-2153(2009)
Sethi, S.K., et al. *Indian Pediatr* 46(5):425-427(2009)