

CDKN2B Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12629a**Specification**

CDKN2B Antibody (N-term) - Product Information

Application	IF, WB, IHC-P, FC,E
Primary Accession	P42772
Other Accession	NP_004927.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	1-30

CDKN2B Antibody (N-term) - Additional Information**Gene ID** 1030**Other Names**

Cyclin-dependent kinase 4 inhibitor B, Multiple tumor suppressor 2, MTS-2, p14-INK4b, p15-INK4b, p15INK4B, CDKN2B, MTS2

Target/Specificity

This CDKN2B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human CDKN2B.

DilutionIF~~1:10~50
WB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CDKN2B Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CDKN2B Antibody (N-term) - Protein Information**Name** CDKN2B

Synonyms MTS2

Function Interacts strongly with CDK4 and CDK6. Potent inhibitor. Potential effector of TGF-beta induced cell cycle arrest.

Cellular Location

Cytoplasm. Note=Also found in the nucleus

Tissue Location


Isoform 2 is expressed in normal (keratinocytes, fibroblasts) and tumor cell lines.

CDKN2B Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CDKN2B Antibody (N-term) - Images




Confocal immunofluorescent analysis of CDKN2B Antibody (N-term)(Cat#AP12629a) with HeLa cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



Western blot analysis of CDKN2B (arrow) using rabbit polyclonal CDKN2B Antibody (N-term) (Cat. #AP12629a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the CDKN2B gene.



CDKN2B Antibody (N-term) (Cat. #AP12629a) immunohistochemistry analysis in formalin fixed and paraffin embedded human lung carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of CDKN2B Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



CDKN2B Antibody (N-term) (Cat. #AP12629a) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

CDKN2B Antibody (N-term) - Background

This gene lies adjacent to the tumor suppressor gene CDKN2A in a region that is frequently mutated and deleted in a wide variety of tumors. This gene encodes a cyclin-dependent kinase inhibitor, which forms a complex with CDK4 or CDK6, and prevents the activation of the CDK kinases, thus the encoded protein functions as a cell growth regulator that controls cell cycle G1 progression. The expression of this gene was found to be dramatically induced by TGF beta, which suggested its role in the TGF beta induced growth inhibition. Two alternatively spliced transcript variants of this gene, which encode distinct proteins, have been reported.

CDKN2B Antibody (N-term) - References

Camacho, C.V., et al. Carcinogenesis 31(10):1889-1896(2010)
Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Pechlivanis, S., et al. Arterioscler. Thromb. Vasc. Biol. 30(9):1867-1872(2010)
Heni, M., et al. Diabetes (2010) In press :
Roder, C., et al. Childs Nerv Syst (2010) In press :