

**K0746 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP12812a**

**Specification**

---

**K0746 Antibody (N-term) - Product Information**

Application	WB, FC,E
Primary Accession	<a href="#">O68CR1</a>
Other Accession	<a href="#">NP_056002.2</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	253-282

**K0746 Antibody (N-term) - Additional Information**

**Gene ID** 23231

**Other Names**

Protein sel-1 homolog 3, Suppressor of lin-12-like protein 3, Sel-1L3, SEL1L3, KIAA0746

**Target/Specificity**

This K0746 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 253-282 amino acids from the N-terminal region of human K0746.

**Dilution**

WB~~1:1000  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

K0746 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**K0746 Antibody (N-term) - Protein Information**

**Name** SEL1L3

**Synonyms** KIAA0746

## Cellular Location


Membrane; Single-pass membrane protein

## K0746 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## K0746 Antibody (N-term) - Images



All lanes : Anti-K0746 Antibody (N-term) at 1:1000 dilution Lane 1: PC-3 whole cell lysate Lane 2: U-251 MG whole cell lysate Lane 3: U-87 MG whole cell lysate Lane 4: A431 whole cell lysate Lane 5: U-2OS whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 125 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



K0746 Antibody (N-term) (Cat. #AP12812a) flow cytometric analysis of U251 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### **K0746 Antibody (N-term) - References**

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :  
Christoforou, A., et al. Mol. Psychiatry 12(11):1011-1025(2007)