

IRAK4 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12875A

Specification

IRAK4 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	O9NWZ3
Other Accession	O8R4K2 , O1RMT8 , NP_001107654.1 , NP_057207.2
Reactivity	Human
Predicted	Bovine, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	25-52

IRAK4 Antibody (N-term) - Additional Information

Gene ID 51135

Other Names

Interleukin-1 receptor-associated kinase 4, IRAK-4, Renal carcinoma antigen NY-REN-64, IRAK4

Target/Specificity

This IRAK4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 25-52 amino acids from the N-terminal region of human IRAK4.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IRAK4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

IRAK4 Antibody (N-term) - Protein Information

Name IRAK4

Function Serine/threonine-protein kinase that plays a critical role in initiating innate immune

response against foreign pathogens. Involved in Toll-like receptor (TLR) and IL-1R signaling pathways (PubMed:[17878374](#)). Is rapidly recruited by MYD88 to the receptor- signaling complex upon TLR activation to form the Myddosome together with IRAK2. Phosphorylates initially IRAK1, thus stimulating the kinase activity and intensive autophosphorylation of IRAK1. Phosphorylates E3 ubiquitin ligases Pellino proteins (PELI1, PELI2 and PELI3) to promote pellino-mediated polyubiquitination of IRAK1. Then, the ubiquitin- binding domain of IKBKG/NEMO binds to polyubiquitinated IRAK1 bringing together the IRAK1-MAP3K7/TAK1-TRAF6 complex and the NEMO-IKKA-IKKB complex. In turn, MAP3K7/TAK1 activates IKKs (CHUK/IKKA and IKBKB/IKKB) leading to NF-kappa-B nuclear translocation and activation. Alternatively, phosphorylates TIRAP to promote its ubiquitination and subsequent degradation. Phosphorylates NCF1 and regulates NADPH oxidase activation after LPS stimulation suggesting a similar mechanism during microbial infections.

Cellular Location


Cytoplasm.

IRAK4 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IRAK4 Antibody (N-term) - Images



IRAK4 Antibody (N-term) (Cat. #AP12875a) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the IRAK4 antibody detected the IRAK4 protein (arrow).



Anti-IRAK4 Antibody (N-term) at 1:1000 dilution + K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 52 kDa Blocking/Dilution buffer: 5% NFD/MBST.

IRAK4 Antibody (N-term) - Background

This gene encodes a kinase that activates NF-kappaB in both the Toll-like receptor (TLR) and T-cell receptor (TCR) signaling pathways. The protein is essential for most innate immune responses. Mutations in this gene result in IRAK4 deficiency and recurrent invasive pneumococcal disease. Multiple transcript variants encoding different isoforms have been found for this gene.

IRAK4 Antibody (N-term) - References

Silva, L.K., et al. Eur. J. Hum. Genet. 18(11):1221-1227(2010)
Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
McDonald, D.R., et al. J. Allergy Clin. Immunol. 126(2):332-337(2010)
Schuurhof, A., et al. Pediatr. Pulmonol. 45(6):608-613(2010)
Wang, Z., et al. Structure 14(12):1835-1844(2006)

IRAK4 Antibody (N-term) - Citations

- [The anti-inflammatory effect and potential mechanism of cardamomin in DSS-induced colitis.](#)