

GAGE12H Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13637a

Specification

GAGE12H Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	A6NDE8
Other Accession	O9UEU5 , A6NER3 , POCL82 , POCL81 , POCL80 , A1L429 , O76087 , Q13070 , Q13069 , Q13068 , Q13067 , Q13065 , Q4V326 , Q13066 , Q6NT46 , Q4V321 , A6NGK3 , NP_001091880.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	12924
Antigen Region	21-49

GAGE12H Antibody (N-term) - Additional Information

Gene ID 729442

Other Names

G antigen 12H, GAGE-12H, GAGE12H

Target/Specificity

This GAGE12H antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 21-49 amino acids from the N-terminal region of human GAGE12H.

Dilution

WB~~1:1000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GAGE12H Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

GAGE12H Antibody (N-term) - Protein Information


Name GAGE12H

GAGE12H Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GAGE12H Antibody (N-term) - Images



GAGE12H Antibody (N-term) (Cat. #AP13637a) western blot analysis in ZR-75-1 cell line lysates (35ug/lane). This demonstrates the GAGE12H antibody detected the GAGE12H protein (arrow).



GAGE12H Antibody (N-term) (Cat. #AP13637a) immunohistochemistry analysis in formalin fixed and paraffin embedded human testis tissue followed by peroxidase conjugation of the secondary

antibody and DAB staining. This data demonstrates the use of GAGE12H Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

GAGE12H Antibody (N-term) - Background

The function of this protein remains unknown.

GAGE12H Antibody (N-term) - References

Gjerstorff, M.F., et al. Tissue Antigens 71(3):187-192(2008)
Ross, M.T., et al. Nature 434(7031):325-337(2005)