

**ANP32A Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP13728c****Specification**

---

**ANP32A Antibody (Center) - Product Information**

Application	<b>WB, IHC-P,E</b>
Primary Accession	<a href="#">P39687</a>
Other Accession	<a href="#">O6PAF6</a> , <a href="#">P51122</a> , <a href="#">NP_006296.1</a>
Reactivity	<b>Human</b>
Predicted	<b>Bovine, Xenopus</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>28585</b>
Antigen Region	<b>133-162</b>

**ANP32A Antibody (Center) - Additional Information****Gene ID** 8125**Other Names**

Acidic leucine-rich nuclear phosphoprotein 32 family member A, Acidic nuclear phosphoprotein pp32, pp32, Leucine-rich acidic nuclear protein, LANP, Mapmodulin, Potent heat-stable protein phosphatase 2A inhibitor I1PP2A, Putative HLA-DR-associated protein I, PHAPI, ANP32A, C15orf1, LANP, MAPM, PHAP1

**Target/Specificity**

This ANP32A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 133-162 amino acids from the Central region of human ANP32A.

**Dilution**

WB~~1:2000  
IHC-P~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ANP32A Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**ANP32A Antibody (Center) - Protein Information**

**Name** ANP32A

**Synonyms** C15orf1, LANP, MAPM, PHAP1

**Function** Multifunctional protein that is involved in the regulation of many processes including tumor suppression, apoptosis, cell cycle progression or transcription (PubMed:[10400610](#), PubMed:[11360199](#), PubMed:[16341127](#), PubMed:[18439902](#)). Promotes apoptosis by favouring the activation of caspase-9/CASP9 and allowing apoptosome formation (PubMed:[18439902](#)). In addition, plays a role in the modulation of histone acetylation and transcription as part of the INHAT (inhibitor of histone acetyltransferases) complex. Inhibits the histone- acetyltransferase activity of EP300/CREBBP (CREB-binding protein) and EP300/CREBBP-associated factor by histone masking (PubMed:[11830591](#)). Preferentially binds to unmodified histone H3 and sterically inhibiting its acetylation and phosphorylation leading to cell growth inhibition (PubMed:[16341127](#)). Participates in other biochemical processes such as regulation of mRNA nuclear-to-cytoplasmic translocation and stability by its association with ELAVL1 (Hu-antigen R) (PubMed:[18180367](#)). Plays a role in E4F1-mediated transcriptional repression as well as inhibition of protein phosphatase 2A (PubMed:[15642345](#), PubMed:[17557114](#)).

#### **Cellular Location**

Nucleus. Cytoplasm Endoplasmic reticulum. Note=Translocates to the cytoplasm during the process of neuritogenesis (By similarity). Shuttles between nucleus and cytoplasm. {ECO:0000250, ECO:0000269|PubMed:18180367}

#### **Tissue Location**


Expressed in all tissues tested. Highly expressed in kidney and skeletal muscle, moderate levels of expression in brain, placenta and pancreas, and weakly expressed in lung. Found in all regions of the brain examined (amygdala, caudate nucleus, corpus callosum, hippocampus and thalamus), with highest levels in amygdala


#### **ANP32A Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **ANP32A Antibody (Center) - Images**





All lanes : Anti-ANP32A Antibody (Center) at 1:2000 dilution Lane 1: Jurkat whole cell lysate Lane 2: HeLa whole cell lysate Lane 3: Raji whole cell lysate Lane 4: Daudi whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 29 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



ANP32A Antibody (Center) (Cat. #AP13728c) immunohistochemistry analysis in formalin fixed and paraffin embedded human tonsil tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of ANP32A Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

### ANP32A Antibody (Center) - Background

Implicated in a number of cellular processes, including proliferation, differentiation, caspase-dependent and caspase-independent apoptosis, suppression of transformation (tumor suppressor), inhibition of protein phosphatase 2A, regulation of mRNA trafficking and stability in association with ELAVL1, and inhibition of acetyltransferases as part of the INHAT (inhibitor of histone acetyltransferases) complex. Plays a role in E4F1-mediated transcriptional repression.

### ANP32A Antibody (Center) - References

- Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
- Habrukowich, C., et al. J. Biol. Chem. 285(35):26825-26831(2010)
- Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)

Valdes, A.M., et al. Arthritis Rheum. 60(7):2046-2054(2009)  
Matilla, A., et al. Cerebellum 4(1):7-18(2005)