

**Mouse Camk2d Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP13799c**

## Specification

---

### Mouse Camk2d Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	<a href="#">Q6PHZ2</a>
Other Accession	<a href="#">Q9DG02</a> , <a href="#">P15791</a> , <a href="#">O77708</a> , <a href="#">Q95266</a> , <a href="#">Q13557</a> , <a href="#">Q5ZKI0</a> , <a href="#">Q2HJF7</a> , <a href="#">P11275</a> , <a href="#">P11798</a> , <a href="#">Q9UQM7</a> , <a href="#">Q6DGS3</a> , <a href="#">Q6DEH3</a> , <a href="#">NP_001020609.1</a> , <a href="#">NP_076302.1</a>
Reactivity Predicted	Human, Mouse Zebrafish, Rat, Bovine, Chicken, Pig, Rabbit, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	56369
Antigen Region	221-250

### Mouse Camk2d Antibody (Center) - Additional Information

**Gene ID** 108058

#### Other Names

Calcium/calmodulin-dependent protein kinase type II subunit delta, CaM kinase II subunit delta, CaMK-II subunit delta, Camk2d, Kiaa4163

#### Target/Specificity

This Mouse Camk2d antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 221-250 amino acids from the Central region of mouse Camk2d.

#### Dilution

WB~~1:1000  
IHC-P~~1:10~50

#### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

Mouse Camk2d Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Mouse Camk2d Antibody (Center) - Protein Information

**Name** Camk2d

**Synonyms** Kiaa4163

**Function** Calcium/calmodulin-dependent protein kinase involved in the regulation of Ca(2+) homeostasis and excitation-contraction coupling (ECC) in heart by targeting ion channels, transporters and accessory proteins involved in Ca(2+) influx into the myocyte, Ca(2+) release from the sarcoplasmic reticulum (SR), SR Ca(2+) uptake and Na(+) and K(+) channel transport (PubMed:[12676814](#), PubMed:[15456698](#), PubMed:[17124532](#)). Targets also transcription factors and signaling molecules to regulate heart function. In its activated form, is involved in the pathogenesis of dilated cardiomyopathy and heart failure (PubMed:[12676814](#), PubMed:[19179290](#), PubMed:[19381018](#)). Contributes to cardiac decompensation and heart failure by regulating SR Ca(2+) release via direct phosphorylation of RYR2 Ca(2+) channel on 'Ser-2808' (PubMed:[20194790](#)). In the nucleus, phosphorylates the MEF2 repressor HDAC4, promoting its nuclear export and binding to 14-3-3 protein, and expression of MEF2 and genes involved in the hypertrophic program. Is essential for left ventricular remodeling responses to myocardial infarction (PubMed:[15793582](#)). In pathological myocardial remodeling acts downstream of the beta adrenergic receptor signaling cascade to regulate key proteins involved in ECC. Regulates Ca(2+) influx to myocytes by binding and phosphorylating the L-type Ca(2+) channel subunit beta-2 CACNB2. In addition to Ca(2+) channels, can target and regulate the cardiac sarcolemmal Na(+) channel Nav1.5/SCN5A and the K+ channel Kv4.3/KCND3, which contribute to arrhythmogenesis in heart failure (PubMed:[17124532](#)). Phosphorylates phospholamban (PLN/PLB), an endogenous inhibitor of SERCA2A/ATP2A2, contributing to the enhancement of SR Ca(2+) uptake that may be important in frequency- dependent acceleration of relaxation (FDAR) and maintenance of contractile function during acidosis. May participate in the modulation of skeletal muscle function in response to exercise, by regulating SR Ca(2+) transport through phosphorylation of PLN/PLB and triadin, a ryanodine receptor-coupling factor. In response to interferon-gamma (IFN-gamma) stimulation, catalyzes phosphorylation of STAT1, stimulating the JAK-STAT signaling pathway (PubMed:[11972023](#)).

### Cellular Location

Cell membrane, sarcolemma; Peripheral membrane protein; Cytoplasmic side. Sarcoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side

### Tissue Location


Expressed in cardiac muscle and skeletal muscle. Isoform Delta 2, isoform Delta 6, isoform Delta 6 and isoform Delta 10 are expressed in cardiac muscle. Isoform Delta 2 is expressed in skeletal muscle.

## Mouse Camk2d Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## Mouse Camk2d Antibody (Center) - Images



Western blot analysis of lysate from ZF4 cell line, using (DANRE) camk2d2 Antibody (Cat. #AP13799c). AP13799c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 35ug.




Mouse Camk2d Antibody (Center) (Cat. #AP13799c) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the Camk2d antibody detected the Camk2d protein (arrow).



Western blot analysis of Camk2d (arrow) using rabbit polyclonal Mouse Camk2d Antibody (Center) (Cat. #AP13799c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or

transiently transfected (Lane 2) with the Camk2d gene.



Mouse Camk2d Antibody (Center) (AP13799c) immunohistochemistry analysis in formalin fixed and paraffin embedded mouse heart tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of Mouse Camk2d Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

#### **Mouse Camk2d Antibody (Center) - Background**

CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release (By similarity).

#### **Mouse Camk2d Antibody (Center) - References**

- Martinez-Pena y Valenzuela, I., et al. J. Neurosci. 30(37):12455-12465(2010)
- Toko, H., et al. Circulation 122(9):891-899(2010)
- Kushnir, A., et al. Proc. Natl. Acad. Sci. U.S.A. 107(22):10274-10279(2010)
- Mangmool, S., et al. J. Cell Biol. 189(3):573-587(2010)
- Koval, O.M., et al. Proc. Natl. Acad. Sci. U.S.A. 107(11):4996-5000(2010)