

**ACVR1 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP13856c**

**Specification**

---

**ACVR1 Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q04771</a>
Other Accession	<a href="#">P80201</a> , <a href="#">P37172</a> , <a href="#">Q28041</a> , <a href="#">NP_001104537.1</a> , <a href="#">NP_001096.1</a>
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	57153
Antigen Region	132-161

**ACVR1 Antibody (Center) - Additional Information**

**Gene ID** 90

**Other Names**

Activin receptor type-1, Activin receptor type I, ACTR-I, Activin receptor-like kinase 2, ALK-2, Serine/threonine-protein kinase receptor R1, SKR1, TGF-B superfamily receptor type I, TSR-I, ACVR1, ACVRLK2

**Target/Specificity**

This ACVR1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 132-161 amino acids from the Central region of human ACVR1.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ACVR1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**ACVR1 Antibody (Center) - Protein Information**

**Name** ACVR1

**Synonyms** ACVRLK2

**Function** Bone morphogenetic protein (BMP) type I receptor that is involved in a wide variety of biological processes, including bone, heart, cartilage, nervous, and reproductive system development and regulation (PubMed:[20628059](#), PubMed:[22977237](#)). As a type I receptor, forms heterotetrameric receptor complexes with the type II receptors AMHR2, ACVR2A or ACVR2B (PubMed:[17911401](#)). Upon binding of ligands such as BMP7 or GDF2/BMP9 to the heteromeric complexes, type II receptors transphosphorylate ACVR1 intracellular domain (PubMed:[25354296](#)). In turn, ACVR1 kinase domain is activated and subsequently phosphorylates SMAD1/5/8 proteins that transduce the signal (PubMed:[9748228](#)). In addition to its role in mediating BMP pathway-specific signaling, suppresses TGFbeta/activin pathway signaling by interfering with the binding of activin to its type II receptor (PubMed:[17911401](#)). Besides canonical SMAD signaling, can activate non-canonical pathways such as p38 mitogen-activated protein kinases/MAPKs (By similarity). May promote the expression of HAMP, potentially via its interaction with BMP6 (By similarity).

**Cellular Location**

Membrane; Single-pass type I membrane protein.

**Tissue Location**


Expressed in normal parenchymal cells, endothelial cells, fibroblasts and tumor-derived epithelial cells

**ACVR1 Antibody (Center) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**ACVR1 Antibody (Center) - Images**



Western blot analysis of ACVR1 (arrow) using rabbit polyclonal ACVR1 Antibody (Center) (Cat. #AP13856c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the ACVR1 gene.



Anti-ACVR1 Antibody (Center) at 1:2000 dilution + A549 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 57 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

#### **ACVR1 Antibody (Center) - Background**

Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I ( I and IB) and two type II ( II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. This gene encodes activin A type I receptor which signals a particular transcriptional response in concert with activin type II receptors. Mutations in this gene are associated with fibrodysplasia ossificans progressive.

#### **ACVR1 Antibody (Center) - References**

Canzian, F., et al. Hum. Mol. Genet. 19(19):3873-3884(2010)  
Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)  
Song, G.A., et al. J. Biol. Chem. 285(29):22542-22553(2010)  
Herrera, B., et al. Cancer Res. 69(24):9254-9262(2009)  
Jung, B., et al. PLoS ONE 4 (12), E8308 (2009) :