

**BMP7 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP13859B****Specification**

---

**BMP7 Antibody (C-term) - Product Information**

Application	<b>WB, IHC-P,E</b>
Primary Accession	<a href="#">P18075</a>
Other Accession	<a href="#">NP_001710.1</a>
Reactivity	<b>Human, Mouse</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>49313</b>
Antigen Region	<b>280-309</b>

**BMP7 Antibody (C-term) - Additional Information****Gene ID** 655**Other Names**

Bone morphogenetic protein 7, BMP-7, Osteogenic protein 1, OP-1, Eptotermin alfa, BMP7, OP1

**Target/Specificity**

This BMP7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 280-309 amino acids from the C-terminal region of human BMP7.

**Dilution**WB~~1:1000  
IHC-P~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

BMP7 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**BMP7 Antibody (C-term) - Protein Information****Name** BMP7**Synonyms** OP1

**Function** Growth factor of the TGF-beta superfamily that plays important role in various biological processes, including embryogenesis, hematopoiesis, neurogenesis and skeletal morphogenesis (PubMed:[31208997](#)). Initiates the canonical BMP signaling cascade by associating with type I receptor ACVR1 and type II receptor ACVR2A (PubMed:[12667445](#), PubMed:[9748228](#)). Once all three components are bound together in a complex at the cell surface, ACVR2A phosphorylates and activates ACVR1. In turn, ACVR1 propagates signal by phosphorylating SMAD1/5/8 that travel to the nucleus and act as activators and repressors of transcription of target genes (PubMed:[12478285](#)). For specific functions such as growth cone collapse in developing spinal neurons and chemotaxis of monocytes, uses also BMP2 as type II receptor (PubMed:[31208997](#)). Can also signal through non-canonical pathways such as P38 MAP kinase signaling cascade that promotes brown adipocyte differentiation through activation of target genes, including members of the SOX family of transcription factors (PubMed:[27923061](#)). Promotes the expression of HAMP, this is repressed by its interaction with ERFE (PubMed:[30097509](#)).

#### Cellular Location

Secreted.

#### Tissue Location


Expressed in the kidney and bladder. Lower levels seen in the brain

#### BMP7 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### BMP7 Antibody (C-term) - Images



Western blot analysis of BMP7 (arrow) using rabbit polyclonal BMP7 Antibody (C-term) (Cat. #AP13859b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the BMP7 gene.



BMP7 Antibody (C-term) (Cat. #AP13859b) western blot analysis in mouse heart tissue lysates (35ug/lane). This demonstrates the BMP7 antibody detected the BMP7 protein (arrow).



BMP7 Antibody (C-term) (Cat. #AP13859b) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of BMP7 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

### **BMP7 Antibody (C-term) - Background**

The bone morphogenetic proteins (BMPs) are a family of secreted signaling molecules that can induce ectopic bone growth. Many BMPs are part of the transforming growth factor-beta (TGFB) superfamily. BMPs were originally identified by an ability of demineralized bone extract to induce endochondral osteogenesis in vivo in an extraskeletal site. Based on its expression early in embryogenesis, the BMP encoded by this gene has a proposed role in early development and possible bone inductive activity. [provided by RefSeq].

### **BMP7 Antibody (C-term) - References**

- Hwang, C.J., et al. J Neurosurg Spine 13(4):484-493(2010)
- Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)
- Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)

Jonigk, D., et al. Virchows Arch. 457(3):369-380(2010)

Pegorier, S., et al. Respir. Res. 11, 85 (2010) :

**BMP7 Antibody (C-term) - Citations**

- [Mineralization Effect of Hyaluronan on Dental Pulp Cells via CD44.](#)