

**ACK1 Antibody (Center Y284)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP13936C**

**Specification**

---

**ACK1 Antibody (Center Y284) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q07912</a>
Other Accession	<a href="#">O54967</a> , <a href="#">Q17R13</a> , <a href="#">NP_005772.3</a> , <a href="#">NP_001010938.1</a>
Reactivity	Human
Predicted	Bovine, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	114569
Antigen Region	265-290

**ACK1 Antibody (Center Y284) - Additional Information**

**Gene ID** 10188

**Other Names**

Activated CDC42 kinase 1, ACK-1, Tyrosine kinase non-receptor protein 2, TNK2, ACK1

**Target/Specificity**

This ACK1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 265-290 amino acids from the Central region of human ACK1.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ACK1 Antibody (Center Y284) is for research use only and not for use in diagnostic or therapeutic procedures.

**ACK1 Antibody (Center Y284) - Protein Information**

**Name** TNK2

## Synonyms ACK1

**Function** Non-receptor tyrosine-protein and serine/threonine-protein kinase that is implicated in cell spreading and migration, cell survival, cell growth and proliferation. Transduces extracellular signals to cytosolic and nuclear effectors. Phosphorylates AKT1, AR, MCF2, WASL and WWOX. Implicated in trafficking and clathrin-mediated endocytosis through binding to epidermal growth factor receptor (EGFR) and clathrin. Binds to both poly- and mono-ubiquitin and regulates ligand-induced degradation of EGFR, thereby contributing to the accumulation of EGFR at the limiting membrane of early endosomes. Downstream effector of CDC42 which mediates CDC42-dependent cell migration via phosphorylation of BCAR1. May be involved both in adult synaptic function and plasticity and in brain development. Activates AKT1 by phosphorylating it on 'Tyr-176'. Phosphorylates AR on 'Tyr-267' and 'Tyr-363' thereby promoting its recruitment to androgen-responsive enhancers (AREs). Phosphorylates WWOX on 'Tyr-287'. Phosphorylates MCF2, thereby enhancing its activity as a guanine nucleotide exchange factor (GEF) toward Rho family proteins. Contributes to the control of AXL receptor levels. Confers metastatic properties on cancer cells and promotes tumor growth by negatively regulating tumor suppressor such as WWOX and positively regulating pro-survival factors such as AKT1 and AR. Phosphorylates WASP (PubMed:[20110370](#)).

## Cellular Location

Cell membrane. Nucleus. Endosome {ECO:0000250|UniProtKB:O54967} Cell junction, adherens junction. Cytoplasmic vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic vesicle, clathrin-coated vesicle Membrane, clathrin-coated pit. Cytoplasm, perinuclear region. Cytoplasm, cytosol {ECO:0000250|UniProtKB:O54967}. Note=The Tyr-284 phosphorylated form is found both in the membrane and nucleus (By similarity). Co-localizes with EGFR on endosomes (PubMed:20333297). Nuclear translocation is CDC42-dependent (By similarity). Detected in long filamentous cytosolic structures where it co-localizes with CTPS1 (By similarity) {ECO:0000250|UniProtKB:O54967, ECO:0000269|PubMed:20333297}

## Tissue Location


The Tyr-284 phosphorylated form shows a significant increase in expression in breast cancers during the progressive stages i.e. normal to hyperplasia (ADH), ductal carcinoma in situ (DCIS), invasive ductal carcinoma (IDC) and lymph node metastatic (LNMM) stages. It also shows a significant increase in expression in prostate cancers during the progressive stages.


## ACK1 Antibody (Center Y284) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## ACK1 Antibody (Center Y284) - Images





ACK1 Antibody (Center Y284) (Cat. #AP13936c) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the ACK1 antibody detected the ACK1 protein (arrow).

#### **ACK1 Antibody (Center Y284) - Background**

This gene encodes a tyrosine kinase that binds Cdc42Hs in its GTP-bound form and inhibits both the intrinsic and GTPase-activating protein (GAP)-stimulated GTPase activity of Cdc42Hs. This binding is mediated by a unique sequence of 47 amino acids C-terminal to an SH3 domain. The protein may be involved in a regulatory mechanism that sustains the GTP-bound active form of Cdc42Hs and which is directly linked to a tyrosine phosphorylation signal transduction pathway. Several alternatively spliced transcript variants have been identified from this gene, but the full-length nature of only two transcript variants has been determined.

#### **ACK1 Antibody (Center Y284) - References**

- Mahajan, K., et al. J. Cell. Physiol. 224(2):327-333(2010)
- Chua, B.T., et al. Mol Oncol 4(4):323-334(2010)
- Prieto-Echague, V., et al. J. Biol. Chem. 285(14):10605-10615(2010)
- Lin, Q., et al. Mol. Cell. Biol. 30(6):1541-1554(2010)
- Pao-Chun, L., et al. J. Biol. Chem. 284(50):34954-34963(2009)