

**PDCD4 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP14001c**

**Specification**

---

**PDCD4 Antibody (Center) - Product Information**

Application	WB, IHC-P-Leica,E
Primary Accession	<a href="#">Q53EL6</a>
Other Accession	<a href="#">Q9JID1</a> , <a href="#">Q61823</a> , <a href="#">NP_055271.2</a> , <a href="#">NP_663314.1</a>
Reactivity	Human, Mouse, Rat
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	121-149

**PDCD4 Antibody (Center) - Additional Information**

**Gene ID** 27250

**Other Names**

Programmed cell death protein 4, Neoplastic transformation inhibitor protein, Nuclear antigen H731-like, Protein 197/15a, PDCD4, H731

**Target/Specificity**

This PDCD4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 121-149 amino acids from the Central region of human PDCD4.

**Dilution**

WB~~1:2000  
IHC-P-Leica~~1:500

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PDCD4 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**PDCD4 Antibody (Center) - Protein Information**

**Name** PDCD4

## Synonyms H731

**Function** Inhibits translation initiation and cap-dependent translation. May exert its function by hindering the interaction between EIF4A1 and EIF4G. Inhibits the helicase activity of EIF4A. Modulates the activation of JUN kinase. Down-regulates the expression of MAP4K1, thus inhibiting events important in driving invasion, namely, MAPK85 activation and consequent JUN-dependent transcription. May play a role in apoptosis. Tumor suppressor. Inhibits tumor promoter-induced neoplastic transformation. Binds RNA (By similarity).

## Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q61823}. Cytoplasm {ECO:0000250|UniProtKB:Q61823}. Note=Shuttles between the nucleus and cytoplasm (By similarity). Predominantly nuclear under normal growth conditions, and when phosphorylated at Ser-457 (PubMed:16357133)

## Tissue Location


Up-regulated in proliferative cells. Highly expressed in epithelial cells of the mammary gland. Reduced expression in lung cancer and colon carcinoma.

## PDCD4 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## PDCD4 Antibody (Center) - Images




All lanes : Anti-PDCD4 Antibody (Center) at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: 293 whole cell lysate Lane 5: HL-60 whole cell lysate Lane 6: Mouse thymus tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band


size : 52 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



All lanes : Anti-PDCD4 Antibody (Center) at 1:2000 dilution Lane 1: Hela whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: Mouse thymus tissue lysate Lane 5: Rat thymus tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 52 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



Immunohistochemical analysis of AP14001C on paraffin-embedded human thyroid tissue was performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:500) for 15min at room temperature. Leica Bond Polymer Refine Detection was used as the secondary antibody.



Immunohistochemical analysis of AP14001C on paraffin-embedded human testis tissue was performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:500) for 15min at room temperature. Leica Bond Polymer Refine Detection was used as the secondary antibody.

#### **PDCD4 Antibody (Center) - Background**

This gene encodes a protein localized to the nucleus in proliferating cells. Expression of this gene is modulated by cytokines in natural killer and T cells. The gene product is thought to play a role in apoptosis but the specific role has not yet been determined. Two transcripts encoding different isoforms have been identified.

#### **PDCD4 Antibody (Center) - References**

Hayashi, A., et al. Hum. Pathol. 41(11):1507-1515(2010)  
Zhang, X., et al. Cancer Sci. 101(10):2163-2170(2010)  
Wang, W., et al. Exp. Cell Res. 316(15):2456-2464(2010)  
Santhanam, A.N., et al. Oncogene 29(27):3921-3932(2010)  
Fassan, M., et al. Oncol. Rep. 24(1):135-139(2010)